

DOCUMENT SOURCE
Lawrence Berkeley Laboratory
Archives and Records Office
SCIENTISTS' PAPERS
JOHN LAWRENCE

Records Series Title _____

Accession No. 434-92-0016
 File Code No. 14-14-0
 Carton No. 2
 Folder No. EUGENE LISBENDER
 Notes INFO ON BONE SEEKING
 Found By KAREN THOMAS
 Date: _____

COPY

724875

AGENTS & BONE IMAGING. INCLUDES INFO ON OLDER "OBSOLETE"
BONE SEEKING AGENTS

I. L. Schaper, M.D.
 M. Fernandez, M.D.
 E.B. Silberstein, M.D.
 R. David, M.D.
 J. Graham, D.O.
 Radioisotope Laboratory
 University of Cincinnati College of Medicine
 1981

File under Eugene Scars
Prophylaxis
Special material

Need to Swift 1st
Cannot read scans
as one needs X-ray
help to complete
Need for quality scans
Pinhole + converging
collimators




BONE IMAGING

I. RADIOPHARMACEUTICALSA. ^{99m}Tc LABELLED PHOSPHATE COMPOUNDS

These include a number of compounds but the following preparations
are most frequently used:

1. ^{99m}Tc-Ethanehydroxy Diphosphonate (^{99m}TcEHDP)
2. ^{99m}Tc-Polyphosphate (^{99m}TcPoly)
3. ^{99m}Tc-Pyrophosphate (^{99m}TcPyro)
4. ^{99m}Tc-Methylene Diphosphonate (^{99m}TcMDP)
5. ^{99m}Tc-Hydroxymethylene Diphosphonate (^{99m}TcHMDP)

Done



ETHANE-1-HYDROX

BEST COPY AVAILABLE

1173218

DOCUMENT SOURCE
Lawrence Berkeley Laboratory
Archives and Records Office
SCIENTISTS' PAPERS

Records Series Title JOHN LAWRENCE

Accession No. 434-92-2216
File Code No. 18-112-6
Carton No. 2
Folder No. EUGENE L. SAENGER
Notes INFO. ON BONE SEEKING
Found By KAREN HOLMES
Date:

COPY

AGENTS & BONE IMAGING. INCLUDES INFO ON OLDER "OBSOLETE" BONE SEEKING AGENTS

E.L. Saenger, M.D.
M. Fernandez, M.D.
E.B. Silberstein, M.D.
R. David, M.D.
J. Graham, D.O.
Radioisotope Laboratory
University of Cincinnati College of Medicine
1981

*File under Eugene Saenger
Phosphates*

*Need to Supt 1st
Cannot read scan
as one needs X-Ray
info to complete
Need for quality scans
Pulse + conversion
collimators*

BONE IMAGING

I. RADIOPHARMACEUTICALS

A. ^{99m}Tc LABELLED PHOSPHATE COMPOUNDS

These include a number of compounds but the following preparations

are most frequently used:

1. ^{99m}Tc-Ethanedihydroxy Diphosphonate (^{99m}TcEDHP)
2. ^{99m}Tc-Polyphosphate (^{99m}TcPoly)
3. ^{99m}Tc-Pyrophosphate (^{99m}TcPyro)
4. ^{99m}Tc-Methylene Diphosphonate (^{99m}TcMDP)
5. ^{99m}Tc-Hydroxymethylene Diphosphonate (^{99m}TcHMDP)

Done




FIG. 1 Chemical formulas of bone seeking phosphates and diphosphonates forming complexes of ^{99m}Tc. Although these are shown as tetrasodium salts, in solution at neutral pH they are probably disodium salts.

These five preparations have been shown to have biologic differences apparently related to their chemical structures. The mechanism of bone uptake of these compounds is not perfectly elucidated but they seem to bind strongly to the bone surface by absorption ("chemisorption"). A recent report suggests that these compounds, and especially pyrophosphate, have a greater affinity for non-osteoid organic matrix than the crystal surface.

DOCUMENT SOURCE Lawrence Berkeley Laboratory Archives and Records Office <i>SCIENTISTS' PAPERS</i>	
Records Series Title	<i>JOHN LAWRENCE</i>
Accession No.	<i>434-92-0016</i>
File Code No.	<i>18-14-0</i>
Carton No.	<i>2</i>
Folder No.	<i>SAENGER, EUGENE L.</i>
Notes	
Found By	<i>Karen Holmes</i>
Date	

COPY

BUSI. IMAGI.GC

All five agents have a high affinity for bone but they also have biological differences which seem to be related to their differences in chemical structure (4,5).

The blood clearance of the Tc-Phosphates in humans and animals appears to be best for the MDP and HMDP, followed by EMDP, Pyrophosphate, and Polyphosphate.




FIG. 2 Blood clearance of MDP in humans compared with three other ^{99m}Tc complexes and ⁹⁹Tc (corrected for physical decay), assuming blood volume was 7% of body weight. PPI indicates pyrophosphate and PolyP denotes polyphosphate.

Subramanian et al. J Nuc Med 16: 748, 1975.

The relatively slower clearance of the ^{99m}Tc-Pyrophosphate and ^{99m}Tc-Polyphosphate is explained by the stronger affinity and binding to serum proteins.

*? Also for
various
isotopes
of...*

1173220

DOCUMENT SOURCE Lawrence Berkeley Laboratory Archives and Records Office SCIENTISTS' PAPERS	
Records Series Title	JOHN LAWRENCE
Accession No.	424-92-0216
File Code No.	19-14-0
Carton No.	(2)
Folder No.	
Notes	
Found By	Karen Holmes
Date	

COPY

BONE IMAGING

3

Overall, Tc-diphosphonate has been favored as the best agent for clinical use followed by the Tc-pyrophosphate and the polyphosphates (10,11).

As an example, some specific data are given for the labelled diphosphonate.

B. TECHNETIUM LABELLED DIPHOSPHONATES (P-C-P Bonds [Pyrophosphonates have P-O-P bonds which lead to some minor differences]).

1. Actions

- a. Increased blood flow
- b. Chemisorption
- c. Displacement of orthophosphate

2. Localization

- a. Amount and type of calcium and phosphorus present
- b. Increased vascularity of area (i.e., epiphyses)
- c. Increased surface area
- d. Increased turnover

3. Physiology and Pharmacology

- a. Optimum scan time - 2-3 hours, depending on the pharmaceutical 50% in bone - 50% excreted. Patient should void at 1 hour and at time of imaging.
- b. Diphosphonate - chelate excreted via GFR (bladder is critical organ)
- c. High ratio bone/soft tissue
- d. High ratio lesion to normal bone
- e. Low radiation dose

1173221

DOCUMENT SOURCE Lawrence Berkeley Laboratory Archives and Records Office SCIENTISTS' PAPERS	
Records Series Title	JOHN LAWRENCE
Accession No.	434-72-2016
File Code No.	18-14-0
Carton No.	2
Folder No.	
Notes	
Found By	Karen Holmes
Date	

COPY

TABLE I

DOSIMETRY OF BONE IMAGING AGENTS
(in 70 kg Adult - Rads/mci Administered) (Data from package insert)

ORGAN	Radiopharmaceutical				
	^{99m} TcMDP	^{99m} Tc Pyrophosphate	^{99m} Tc EHDP	¹⁸ F	⁶⁷ Ga Citrate
Skeleton	.035	.034	.039	.15-.29	.44
Bone Marrow	.028	.028-.036	.028	.04	.58
Kidneys	.031	.028-.14	.14		.41
Liver	.008	.001	----		.46
Total Body	.0065	.009	.009	.05	.26
Bladder				2.0	
2-3 h voiding	.13	.04-.1	.1		----
4.8-6 h voiding	.31	.07-.23	.23		----
Testis					.24
2-3 h voiding	.008	.004-.01			
4.8-6 h voiding	.01	.007-.015			
Ovaries					.28
2-3 h voiding	.012	.004-.009			
4.8-6 h voiding	.017	.007-.015			
Heart					----
Normal		.007	.007		
Impaired		.014	.02-.03		
Gastroenteric Tract					
Stomach					.22
Small bowel					.36
Upper large bowel					.56
Lower large bowel					.9

*Doses 10-20 mCi
Blood pool sep infection
Average 3-5 hrs, 24 hrs.
3.5 mCi/mCi of administered activity to bone
kidneys dose 2x
marrow " 1/2
TB dose 10 mCi/mCi
Bladder wall > 10 rads/study
hydrate + void*

1173222

DOCUMENT SOURCE Lawrence Berkeley Laboratory Archives and Records Office SCIENTISTS' PAPERS	
Records Series Title	JOHN LAWRENCE
Accession No.	434-92-0016
File Code No.	14-14-6
Carton No.	2
Folder No.	
Notes	
Found By	Karen Holmes
Date:	

COPY

BONE IMAGING

C. OLDER BONE SEEKING AGENTS

1. ^{18}F - short T_{1/2} (1.8 hr). High energy (511 KeV) annihilation photons are not ideal for gamma camera.

^{18}F has very favorable biologic characteristics. It is absorbed by the GI tract and can be given orally. The mechanism of uptake appears to be that of ion exchange at the level of the hydroxyapatite crystals. Extraction efficiency is excellent and approaches 100% depending on bone flow.

^{18}F is rapidly secreted by kidney (20-60% at 2 hours) a factor to keep in mind since overlapping of the kidneys and bladder with structures may occur.

2. ^{85}Sr - long T_{1/2} (64 days) requires low administered dose for low patient doses. This leads to low photon flux, long scanning time, and low resolution. Obsolete.

3. ^{87}Sr - short T_{1/2} (2.8 hours) does not allow for adequate soft tissue clearance. Higher photon energy (388 KeV) not ideal for gamma cameras. Obsolete.

DOCUMENT SOURCE Lawrence Berkeley Laboratory Archive and Records Office SCIENTISTS' PAPERS	
Records Series Title	JOHN LAWRENCE
Accession No.	432-92-0016
File Code No.	18-14-6
Carton No.	2
Folder No.	
Notes	
Found By	Karen Holmes
Date:	

COPY

BONE IMAGING

C. The Thoracic Region

1. Inferior angle and spine of scapula are well seen. Sternum and ossification centers and sternoclavicular joints (watch for asymmetry. Occasionally, costo chondral junctions are well seen.

D. The Long Bones

1. Small bones in wrists, hands, and feet are frequently easily visible due to large proportion of cancellous to cortical bones. Good detail achieved with pinhole collimator.

E. Other Normal Sites of Increased Uptake

1. Epiphyses and apophyses in juveniles.
2. Joints - variable and sometimes asymmetric.
 - a. Sacro-iliac joints usually show increased uptake. Sacral tubercle normally prominent (~4%). [Blei L., et al., The Sacral Tubercle - A Cause for Hot Spots on Bone Scan. Clin Nuc Med 3: 351-354, 1978].
 - b. Skull - petrous pyramids and facial bones
 - c. Cartilage - watch for thyroid cartilage
 - d. Shoulder uptake corresponding to handedness and/or position.
3. Organs
 - a. Kidney and bladder
 - b. Breast
 - c. Myocardium
4. Free $^{99m}\text{TcO}_4$
 - a. Thyroid
 - b. Salivary Gland, Sinuses
 - c. Stomach
 - d. Bowel
 - e. Liver (rare)
5. Technical - change in distance of spine from collimator

* IV. INTERPRETATION - ABNORMAL

- * A. Non-specificity of Bone Scans - cannot distinguish benign from malignant disease.
- B. Soft Tissue Injury
Post-operative sites, myocardial infarction, CVA.
- C. Vertebral Bodies
Most common sites for bony metastasis.
- D. Delayed Scanning
 - * 1. In patients who cannot empty bladder; to increase target/non-target ratio.

1173225

DOCUMENT SOURCE Lawrence Berkeley Laboratory Archives and Records Office SCIENTISTS' PAPERS	
Records Series Title	JOHN LAWRENCE
Accession No.	432-92-00161
File Code No.	19-14-0
Carton No.	(2)
Folder No.	
Notes	
Found By	Karen Holmes
Date	

COPY

BONE IMAGING

D. Delayed Scanning (continued)

2. In patients with questionable densities at 2-3 hours. At 12-24 hours occasionally an abnormality will appear better visualized than at the earlier time.

8
Voeding

E. Spot Images using pinhole or other collimator (converging or diverging) as indicated is often useful. Oblique and lateral views are often helpful. The opposite side should be imaged for comparison. *

F. Metastases

Metastases to the skeleton may show as single or multiple areas of increased uptake. Practically any bone of the skeleton may be involved, but spine, ribs, and pelvis are the most frequently observed.

The most frequent tumors with bone metastases found in clinical practice are cancer of the breast, lung and prostate gland (12).

A clear decision for preoperative bone scans in patients with operable Breast Cancer is not entirely resolved. Patients with lesions < 2 cm in diameter tend not to have positive scans. With increase in lesion size and the presence of positive axillary nodes rates of true positive scans range from 6% to about 25%. The variation is due to several factors: size of tumor, differences in hospital populations, differences in technique, problems of FP and FN interpretation. FN include early very small metastases, lack of an osteoblastic response, lesions of pubis and ischium. FP include the many other causes of positive bone scans. CT, observation and biopsy may be needed to resolve this problem. The subsequent development of metastases is related to the size and extent of the original tumor.

References: Bone scanning in breast cancer, Brit. M.J. 2: 180-181, 15 July 1978.

Galasko CSB. Problems associated with the detection of skeletal metastases. J. Roy. Soc. Med. 71: 38-41, 1978.

McNeil BJ et al. Preoperative and follow up of bone scans in patients with primary carcinoma of the breast. Surg, Gynec and Obst 147: 745-748, 1978.

Clark DG et al. Indications for bone scans in preoperative evaluation of breast cancer. Am J Surg 135: 667-670, 1978.

G. The Patient With Single Bone Lesions

Single bone lesions in bone scanning are frequently found (13). The most common causes are:

1. Metastatic tumor (lung, breast, prostate)
2. Post thoracotomy (surgery scar)
3. Fracture of vertebral body
4. Normal variants (shoulder, sternum)

1173226

DOCUMENT SOURCE	
Lawrence Berkeley Laboratory Archives and Records Office	
SCIENTISTS' PAPERS: JOHN LAWRENCE	
Records Series Title	
Accession No.	42-72-0066
File Code No.	12-14-0
Carton No.	(2)
Folder No.	
Notes	
Found By	Karen Holmes
Date	

COPY

BONE IMAGING

H. The Patient with Diffuse Increase Uptake in Skeleton (14,15,16)

Diffuse or symmetric increase uptake is not a rare situation. The importance of this possibility stems from the fact that a symmetric diffuse uptake may be easily overlooked.

This finding has been noted in metastatic cancer of the prostate and diffuse lymphoma and cancer of the renal pelvis. It is also expected in metabolic bone disease due to hyperparathyroidism and osteomalacia. In our lab we have found Paget's of the skull and hypertrophic pulmonary osteoarthropathy present with symmetric bone involvement. It can occur in hypervitaminosis D and also in renal osteodystrophy where it may represent secondary hyperparathyroidism. [Fogelman et al. Bone scan findings in hypervitaminosis D: Case report, J Nuc Med 18: 1205-1207, 1977].

I. Presence of "Cold" Lesions in Bone Scanning (17,18,19)

A cold lesion may occur as the sole abnormality in a bone scan and it has been recognized most frequently in the vertebra and long bones. It has been described in the following conditions:

1. Metastatic disease of different sources
2. Post-traumatic aseptic necrosis
3. Sickle cell anemia with bone infarction
4. Chronic renal failure

Pathogenesis of this type of condition has not been elucidated but interruption of blood supply appears to be a prominent factor. Conceivably a poor osteoblastic reaction surrounding an area of necrosis or tumor would result in a "cold" area.

See Table III.

J. The False Negative and False Positive Bone Scan

False negative bone scans may result when radioactivity in the lesion is not intense enough, when the lesion is mostly osteolytic (e.g., 50% of multiple myeloma) or when the disease process involves the skeleton in a diffuse or symmetric fashion.

False positive scans are seen mainly as a consequence of contamination with urine and at the site of injection. If the technetium label is poor, uptake by the thyroid gland and the stomach may be a source of confusion. Increased uptake in the area of radical mastectomy presumably due to decreased attenuation of γ rays by removed soft tissue has been observed. Increased uptake in areas of post-surgery scars, biopsy sites, normal or pathologic breast tissue, soft tissue, inflammation and myocardial and cerebral infarcts occurs.

K. Increased Concentration in Extraskelatal Tissues

This condition has been found in many circumstances. The causes and conditions can be listed as follows:

1173227

1. Uptake in Osteoid Forming Tissues

DOCUMENT SOURCE Lawrence Berkeley Laboratory Archives and Records Office SCIENTISTS' PAPERS JOHN LAWRENCE	
Records Series Title	
Accession No.	434-92-0210
File Code No.	18-14-6
Carton No.	2
Folder No.	
Notes	
Found By	Karen Holmes
Date	

COPY

BONE IMAGING

10

K. Increased Concentration in Extrasketal Tissues (con't.)

- a. Myositis ossificans
- b. Diffuse calcinosis
- c. Metastatic osteosarcoma
- d. Osseous metaplasia in soft tissues or lymph nodes
- e. Ectopic calcification in paraplegic patients
- f. Metastases of bladder (with osseous metaplasia)
- g. Calcific tendonitis

In the above conditions concentration of bone seeking agents is expected and probably related to the presence of hydroxyapatite as well as immature collagen matrix.

2. "Non-osteoid" forming tissues -

- a. Thoracotomy and post-surgery scars
- b. Soft tissue inflammation
- c. Brain infarction
- d. Brain astrocytoma
- e. Brain metastases from lymphoma
- f. Myocardial infarction

The mechanism of uptake is not known but increased blood flow and concentration of the isotope by the collagen and intracellular structure could be involved.

L. The Significance of Kidney Abnormalities Incidentally Found During Bone Imaging (20-23)

This point has been recently brought to attention since many of the bone seeking radioisotopes, specifically the Tc-Phosphate compounds, are avidly concentrated by the kidneys enabling us to observe some anatomic detail of the kidneys and excretory system. The following points should be emphasized:

1. There is no good correlation between kidney concentration of ^{18}F -Fluoride and renal function.
2. Hence, kidney asymmetry during ^{18}F -Fluoride bone scanning is unreliable to predict renal disease.
3. The overall accuracy and the level of confidence when kidneys are found morphologically abnormal are high.
4. In general, information regarding kidney morphology and function appears to be similar with all four currently used Tc-Phosphate compounds.
5. The most common conditions that can be detected during bone imaging in general clinical practice are -
 - a. Bilateral decreased visualization due to:
 1. bilateral kidney disease
 2. increased bone uptake ("steal phenomena")

1173228

DOCUMENT SOURCE	
Lawrence Berkeley Laboratory Archives and Records Office SCIENTISTS' PAPERS	
Records Series Title	JOHN LAWRENCE
Accession No.	339-92-0016a
File Code No.	19-14-6
Carton No.	(2)
Folder No.	
Notes	
Found By	Karen Holmes
Date	

COPY

BONE IMAGING

11

L. The Significance of Kidney Abnormalities... (con't)

- b. Small kidney due to unilateral disease
- c. Space occupying lesions: tumors, cysts, dilated calyces
- d. Absent kidney: congenitally, surgically, destruction by encroaching process
- e. Obstructive uropathy with dilated excretory system
- f. Displacement of bladder by pelvic tumor
- g. Ectopic location

M. Bone Scanning in Pediatrics

Some pertinent points are to be considered when dealing with the pediatric age group.

1. The presence of normal variants inherent to this group such as the increased uptake in juxta-articular areas.
2. Positive bone scans due to metastatic disease of pediatric neoplasms, i.e. Wilms Tumor, neuroblastoma, leukemia, rhabdomyosarcoma. Valuable in following course of disease and evaluating treatments. In neuroblastoma bone imaging may miss the lesion although x-ray may be positive. [Reference - Kaufman RA et al. False negative bone scans in neuroblastoma metastatic to ends of long bones. Am J Roentgenol. 130: 131-135, 1978].
3. Frequency of solitary lesions are more likely to be associated with primary bone tumors. Benign tumors take up less activity than do malignant ones. Imaging is useful in determining extent of osteoblastic tumors.
4. Differential diagnoses between osteomyelitis and cellulitis. This is a frequent situation which poses a difficult problem since clinical and radiologic findings may not be conclusive. In this regard, comparing early "blood pool" pictures with the conventional delayed scan is often helpful. In soft tissue inflammation or infection, the early blood pool scans show a greater concentration of the radiolotope when compared with the delayed scan. The reverse is usually found in cases of osteomyelitis. High sensitivity and pinhole collimators are often helpful when dealing with small children. Septic arthritis, discitis (spondylarthritis), bone infarcts in sickle cell disease require consideration in the differential diagnosis.
5. Aseptic necrosis. If imaged early, may show decreased uptake. Later uptake will be increased. In Perche's disease osteotomy may decrease uptake. Attention to technique important for bilateral disease and to follow course. Quantitative technique may be valuable.

(References: Norley TR et al. Femoral head activity in

1173229

DOCUMENT SOURCE	
Lawrence Berkeley Laboratory Archives and Records Office <i>SCIENTISTS' PAPERS:</i> <i>JOHN LAWRENCE</i>	
Records Series Title	
Accession No.	<i>43-32-226</i>
File Code No.	<i>14-14-0</i>
Carton No.	<i>2</i>
Folder No.	
Notes	
Found By	<i>Karen Holmes</i>
Date	

COPY

BONE IMAGING

M. Bone Scanning in Pediatrics (con't)

Perthe's disease: Clinical evaluation of a quantitative technique for estimating tracer uptake, J Nucl Med 19: 884-890, 1978.

Sty JR. Panner's disease (Osteonecrosis of the capitellum). Clin Nuc Med 3: 117, 1978].

6. Discitis - Spondylarthritis - Originally described by Saenger (Am J Roentgenol 64: 20-31, 1950) by x-ray. Scanning reveals increased activity in adjacent vertebral bodies. [Reference - Wenger DR et al. The spectrum of intervertebral disc-space infection in children, J Bone & Joint Surg 60A: 100-108, 1978]. In one reported case the TC diphosphonate scan was normal but the ⁶⁷Ga scan was positive. [Reference - Norris S et al. Early diagnosis of disc-space infection using gallium 67. J Nucl Med 19: 384-386, 1978].

7. Metabolic and congenital bone diseases. Activity often increased but not always. Perhaps a function of patient's age. Quantitative techniques may be useful. [Reference - Sty JR et al. Bone scintigraphy demonstrating Englemann's disease. Clin Nuc Med 3: 69-70, 1978].

[References - Whyte MP et al. ^{99m}Tc Pyrophosphate bone imaging in osteopoikilosis, osteopathia striata and melorheostosis. Radiol 127: 439-443, 1978.

Epstein DA et al. Bone scintigraphy in hereditary multiple exostoses. Am J Roentgenol 130: 331-333, 1978.

Park H-M et al. Skeletal and reticuloendothelial imaging in osteopetrosis: Case report. J Nuc Med 18: 1091-1095, 1977].

8. General References - Good review articles

Harccke HJ Jr., Bone imaging in infants and children: A review. J Nuc Med 19: 324-329, 1978.

Conway JJ. Radionuclide bone imaging in pediatrics. Ped Chn N Am 24: 701-712, 1977.

V. COMPARISON OF X-RAYS, EMDP, FLUORINE

Radiology 107: 551-555, June 1973, Silberstein et al.

Ten patients with Ca and suspected metastasis had ^{99m}Tc-EHDP or ¹⁸F bone scans and a roentgenographic skeletal survey.

	¹⁸ F	EMDP	X-RAY	Total Mets
# Mets	33	59	17	60
% of Total	56%	98%	28%	100%

1173230

DOCUMENT SOURCE Lawrence Berkeley Laboratory Archives and Records Office <i>SCIENTISTS' PAPERS</i>	
Records Series Title	<i>JOHN LAWRENCE</i>
Accession No.	<i>439-72-00161a</i>
File Code No.	<i>14-16-10</i>
Carton No.	<i>2</i>
Folder No.	
Notes	
Found By	<i>Karen Holmes</i>
Date	

COPY

BONE IMAGING

VI. GALLIUM 67 FOR BONE IMAGING

^{67}Ga may be useful in circumstances where $^{99\text{m}}\text{Tc}$ phosphorus compounds appear to give normal images as occasionally in osteomyelitis, septic arthritis. In primary bone tumors it correlates well with $^{99\text{m}}\text{Tc}$ compounds. It may be more useful in showing the extent of soft tissue extension. Its chief drawbacks are the high energy of its photons resulting in poor resolution. It is superior to radiography but generally not as reliable as $^{99\text{m}}\text{Tc}$ compounds. In selected cases where $^{99\text{m}}\text{Tc}$ appears equivocal or at variance with expectation it should be utilized. (Reference - Handmaker H et al. Gallium imaging in pediatrics. J Nuc Med 18: 1057-1063, 1977).

VII. DISEASES OF JOINTS

Joint abnormalities frequently show increased activity with $^{99\text{m}}\text{Tc}$ phosphorus compounds. Osteoarthritis joints are frequently visualized in the course of bone surveys. It is prudent to examine these joints clinically or by x-ray to exclude other causes for visualization.

In rheumatoid arthritis bone scanning agents usually show more joints to be involved than are found clinically and other forms of arthritis show similar findings. Septic arthritis is readily visualized. The peri-articular activity is usually increased due to increased blood flow and local reaction. Symmetrical abnormal increased in activity of involved joints is sometimes difficult to resolve especially with involvement of sacro-iliac joints. Although some investigators find obtaining joint to sacrum count ratios to be valuable in finding and staging sacro-iliac disease (Dequeker et al. Scintigraphic investigation of sacro-iliac disease. J Nuc Med 19: 119-120, 1978), and others (Lentle BC et al. J Nuc Med 19: 120, 1978) disagree in its usefulness for the diagnosis of ankylosing spondylitis.

A painful total hip or knee prosthesis suggests possibility of loosening of one or more components. Imaging with a Tc phosphate complex is an effective screening procedure to determine loosening or other complications. The bone scan alone lacks specificity but approaches 100% sensitivity for loosening and or infection. If the bone scan is negative, no further work-up is necessary at that time. Recent interest has been directed to the combined use of Tc-MDP and gallium in the evaluations of a painful total hip prosthesis. Both studies are needed as the interpretation depends on the presence of abnormal uptake plus the pattern of abnormal uptake, that is to assess the congruity or incongruity of the patterns of abnormal uptake.

EVALUATION OF TOTAL JOINT REPLACEMENT WITH Tc-PHOSPHATES AND GALLIUM

	BONE SCAN	GALLIUM SCAN	CONGRUENT	INCONGRUENT
LOOSENING	+	-	-	-
INFECTION	+	+	-	+
OR	+	(INTENSE)	+	-
TRAUMA	+	+	+	-

VIII. QUANTITATIVE TECHNIQUES

Among these methods currently utilized is that using probes (Park et al, A quantitative evaluation of rheumatoid arthritis activity with Tc $^{99\text{m}}$ HEDP, J Nuc Med 18: 973-976, 1977) with counts over affected and normal areas at specific times after.

1170731

DOCUMENT SOURCE	
Lawrence Berkeley Laboratory Archives and Records Office SCIENTISTS' PAPERS	
Records Series Title	JOHN LAWRENCE
Accession No.	434-32-2266
File Code No.	18-14-6
Carton No.	2
Folder No.	LAENGER, EUGENE
Notes	
Found By	Karen Holmes
Date:	

COPY

BONE IMAGING

review of earlier images to evaluate treatment in rheumatoid arthritis. Another method uses regions of interest and external standards to estimate the amounts of activity taken up in various parts of the skeleton in Paget's disease [Lurye DR et al. An improved method for quantitative bone scanning, J Nuc Med 18: 1069-1073, 1977]. A third method uses serial measurements at frequent intervals up to 24 hours with a shadow shield whole body monitor [Fogelman et al. The use of whole body retention of Tc 99m diphosphonate in the diagnosis of metabolic bone disease J Nuc Med 19: 270-275, 1978; see also Holmes RA, editorial, Quantification of skeletal Tc-99m labeled phosphates to detect metabolic bone disease, J Nuc Med 19: 330-331, 1978]. This technique differentiated patients with renal osteodystrophy, Paget's disease, osteomalacia and primary hyperparathyroidism from normal subjects. Patients with osteoporosis did not differ from normals.

IX. MISCELLANEOUS

Careful evaluation of jaws and related structures can reveal positive images associated with extraction sites, pulp and periodontal infections and local irritation [Tow DE et al. Bone scan in dental disease, J Nuc Med 19: 845-847, 1978].

The use of single photon emission tomography of facial bones using a special collimator and Tc Pyrophosphate compounds gives improved resolution of facial bones and improves the study of tumors, infections, bone grafts and post irradiation osteonecrosis [Brown ML et al. Facial bone scanning by emission tomography, J Nuc Med 18: 1184-1188, 1977].

X. GAMUT APPROACH AND REFERENCES

The gamut approach to roentgenology has been applied to nuclear medicine. Though scintiscan abnormalities are less specific than roentgen lesions, the approach has been useful.

A recent gamut prepared by one of our residents is included. It contains a comprehensive list of references.

TABLE II

SINGLE LOCALIZED AREA OF INCREASED RADIONUCLIDE UPTAKE ON A BONE SCAN

COMMON

1. Metastatic tumor (lung, breast, prostate) (26,67)
2. Post thoracotomy/surgery scar (26,28)
3. Vertebral body compression fracture (26)
4. Normal variants (shoulder, sternum) (29)

1173232

DOCUMENT SOURCE	
Lawrence Berkeley Laboratory Archives and Records Office SCIENTISTS' PAPERS	
Records Series Title	JOHN LAWRENCE
Accession No.	432-92-0216
File Code No.	18-14-4
Carton No.	2
Folder No.	
Notes	
Found By	Karen Holmes
Date	

COPY

BONE IMAGING

15

TABLE II (con't)

LESS COMMON

1. Metastatic tumor (cervix, neuroblastoma in children) (27,30)
2. Primary bone tumor (Ewing's sarcoma, osteosarcoma, osteochondroma) (26,31)
3. Lymphoma (26,27)
4. Monarticular degenerative disease (26)
5. Trauma - fracture other than vertebral body, biopsy site, prosthesis site) (26)
6. Osteomyelitis, pyogenic (26)
7. Peridontal disease (33) and post tooth extraction
8. Paget's disease (26,27)

RARE

1. Metastatic tumor (thyroid, renal, melanoma, pancreas, other gastrointestinal) (27)
2. Primary bone tumor (fibrosarcoma, chondrosarcoma, giant cell tumor, fibrous dysplasia, enchondroma, osteoid osteoma, bone cyst, hemangioma) (26, 32-35)
3. Multiple myeloma (27)
4. Aseptic necrosis (32)
5. Osteitis pubis (32)
6. Osteomyelitis, TB (26,36) coccidiomycosis (27)
7. Extra skeletal calcified and non-calcified tissue uptake [breast carcinoma (29,34), neuroblastoma (37,38), neurofibroma (39), brain metastasis from lung carcinoma (34), cecal/rectal carcinoma (34), nasopharyngeal carcinoma (34), fibrosarcoma (34), soft tissue abscess (40), brain infarction (41), myocardial infarction and other areas of tissue necrosis, Hodgkin's involving spleen (42), leukemic infiltrates (35), myositis ossificans (43), calcific tendonitis (32), thrombophlebitis (29)]. Also in lung, stomach, liver, kidneys in metastatic melanoma and chronic renal disease [Veukateah et al. Metastatic calcification: the role of bone scanning, Radiology 129: 755-758, 1978; Rosenthal DI, Uptake of bone imaging agents by diffuse pulmonary metastatic calcification, Am J Roentgenol 129: 871-874, 1977; Oren VO et al., Liver metastases of oat cell carcinoma of lung detected on ^{99m}Tc diphosphonate bone scan, Clin Nuc Med 3: 355-357, 1978 - has a good table of soft tissue uptake with ^{99m}Tc bone imaging agents].

1173233

DOCUMENT SOURCE	
Lawrence Berkeley Laboratory Archives and Records Office SCIENTISTS' PAPERS	
Records Series Title	JOHN LAWRENCE
Accession No.	92-92-0216
File Code No.	19-14-0
Carton No.	②
Folder No.	
Notes	
Found By	KAREN HOLMES
Date	

copy

BONE IMAGING

16

TABLE II (con't)

8. Gout (26)
9. Periostitis (31)
10. Meningiomas (44)
11. Scurvy - [Front D et al., Bone scintigraphy in scurvy, J Nuc Med 19: 916-917, 1978].

TABLE III

DECREASED RADIONUCLIDE UPTAKE ON A BONE SCAN

AVASCULAR AREAS

- Aseptic necrosis, bone infarcts
- Radiation therapy (late effect)
- Lack of weight bearing stress
- Metastasis
- Sickle Cell C crisis
- Chronic renal failure
- Acute osteomyelitis
- References: Georgen et al., J Nuc Med 15: 1120-1124, 1974
- Fordham et al., Sem Nuc Med 4: 411-429, 1974
- Quint PA, Radiol 130: 751-752, 1979, #58.

TABLE IV

True Abnormal, False Abnormal and False Normal Abnormalities in Bone Imaging
Silberstein, EB, J Nuc Med 17: 229-232, 1976

Abnormalities generally show increased uptake of radiopharmaceuticals

- A. True Abnormal
 1. Tumor, primary or secondary
 2. Fractures and surgical osteotomy
 3. "Metabolic"
 - a. hyperparathyroidism
 - b. Paget's disease
 - c. osteoporosis
 - d. osteomalacia and occasional pseudofractures of ribs [Fogelman et al., The role of bone scanning in osteomalacia, J Nuc Med 19: 245-248, 1978; Fogelman et al., Pseudofracture of the ribs detected by bone scanning, J Nuc Med 18: 1236-1237, 1977].
 4. Inflammation of bone
 - a. osteomyelitis
 - b. abscess
 - c. sterile osteitis (e.g. osteitis pubis)
 - d. granuloma including sarcoid, eosinophilic granuloma
 - e. hyperostosis frontalis interna
 - f. fibrous dysplasia
 - g. hypertrophic pulmonary osteoarthropathy

1173234

DOCUMENT SOURCE	
Lawrence Berkeley Laboratory Archives and Records Office SCIENTISTS' PAPERS	
Records Series Title	JOHN LAWRENCE
Accession No.	92A-92-0016a
File Code No.	19-14-0
Carton No.	(2)
Folder No.	
Notes	
Found By	Karen Holmes
Date	

COPY

BONE IMAGING

17

TABLE IV (con't)

5. Arthritis
 - a. osteoarthritis
 - b. rheumatoid arthritis
 - c. gouty arthritis
 6. Soft-tissue calcifications
 - a. myositis ossificans
 - b. soft-tissue osseous metaplasia
 - c. soft-tissue tumors with calcification or ectopic bone formation
 - d. vascular calcification, especially femoral artery
 - e. calcific tendonitis
 - f. abscess
 - g. infarct, cerebral or myocardial
 - h. thrombophlebitis
 7. Vascular
 - a. surrounding the bone infarct
 8. Decreased uptake
 - a. tumor
 - b. disuse of limb (may also be increased with osteoporosis)
 - c. vascular obstruction (e.g., sickle cell disease, aseptic necrosis)
- B. Falsely Abnormal
1. Renal artifacts or disease
 - a. hydrourater-hydronephrosis with ^{99m}Tc-diphosphonate
 - b. contamination of clothing or skin with urine
 2. Recent surgical procedures on bone or soft tissue
 3. Biopsy site
 4. Colloid formation with liver-spleen uptake
- C. Falsely Normal
1. Lesions of smaller size than the resolving power of the system
 2. Purely lytic lesions (e.g., some myelomas)
 3. Jewelry, prostheses, pacemaker overlying a lesion

TABLE V

Bilateral Lower Limb Uptake of Bone Scanning Agents, in
R.P. Spencer and J.A. Datu, Sem Nuc Med 10; (#3): 314-316, 1980

Some causes of bilateral lower limb uptake of bone imaging agents are:

COMMON

1. "Calcified" femoral vessels
2. Trauma (and battered child)
3. Stress fractures
4. Aseptic necrosis,* osteonecrosis

*increased uptake is likely outside of the affected region.

1173235

DOCUMENT SOURCE	
Lawrence Berkeley Laboratory Archives and Records Office	
SCIENTISTS' PAPERS	
Records Series Title	JOHN LAWRENCE
Accession No.	43a-92-0066
File Code No.	14-14-6
Carton No.	2
Folder No.	
Notes	
Found By	Karen Holmes
Date:	

COPY

BONE IMAGING

18

TABLE V (con't).

5. Multiple small vessel occlusion, as in SS disease*
6. Osteomyelitis (acute or chronic)
7. Hypertrophic pulmonary osteoarthropathy
8. Increased collagen turnover**
9. Paget's disease
10. Arthritides (osteoarthritis, rheumatoid)
11. Multiple metastases⁺

UNCOMMON

1. Ectopic calcifications
2. Injection sites
3. Polymyositis, myolysis, dermatomyositis
4. Soft tissue infection, infarction
5. Surgically induced (bone grafts, pinning, prostheses)
6. Septic or other emboli*
7. Bilateral sympathectomy
8. Lymphoma
9. Multiple myeloma
10. Gaucher's disease
11. Sarcoidosis⁺⁺
12. Scurvy
13. Myelofibrosis
14. Joint disorders

*increased uptake is likely outside of the affected region

**Includes hyperthyroidism, acromegaly, thyroid acropachy, hyperparathyroidism, renal osteodystrophy.

⁺Multiple primary bone tumors can occur but are quite rare.

⁺⁺Well documented radiographically, less well established by bone scans.

1173236

DOCUMENT SOURCE Lawrence Berkeley Laboratory Archives and Records Office SCIENTISTS' PAPERS JOHN LAWRENCE	
Records Series Title	
Accession No.	434-72-0016
File Code No.	19-14-26
Carton No.	2
Folder No.	
Notes	
Found By	Karen Holmes
Date	

COPY

REFERENCES

1. Blau, Monte, et al. ¹⁸F-Fluoride for Bone Imaging. Sem Nuc Med Vol 2, January 1972.
2. Moon, N.M. et al. The Clinical Use of Sodium Fluoride F-18 in Bone Photoscanning. JAMA 204: 116-122.
3. O'Mara, R.E. Bone Scanning in Osseous Metastatic Disease. JAMA 229: 1915-1917.
4. Subramanian, G. et al. An Evaluation of ^{99m}Tc-labelled Phosphate Compounds as Bone-Imaging Agents, in Subramanian, G. et al (eds) Radiopharmaceuticals, New York, Society of Nuclear Medicine, Inc., pp 319-328, 1975.
5. Ackerhalt, R.E., et al. A Comparative Study of Three ^{99m}Tc-labelled Phosphorus Compounds and ¹⁸F-Fluoride for Skeletal Imaging. J Nuc Med 15: 1153-1157, 1974.
6. Garcia, Daniel A., et al. Relative Accretion of ^{99m}Tc-Polyphosphate by Forming and Resorbing Bone Systems in Rats: Its Significance in the Pathologic Basis of Bone Scanning, J Nucl Med 17: 93-97, 1976.
7. Bowen, B.M. et al. Analysis of the Relationship between ^{99m}Tc-Sn-Polyphosphate and ^{99m}Tc-Pyrophosphate. J Nucl Med 15: 652-655, 1974.
8. Rosenthal, L. and Kaye, M. ^{99m}Tc-Pyrophosphate Kinetics and Imaging in Metabolic Disease. J Nucl Med 16: 33, 1975.
9. Kaye, M. et al. ^{99m}Tc-Pyrophosphate: Studies in-vivo and in-vitro J Nucl Med 16: 40-45, 1975.
10. Krishnamurphy, G.H. et al. Clinical Comparison of the Kinetics of ^{99m}Tc-Labelled Polyphosphate and Diphosphonate. J Nucl Med 15: 848-855, 1974.
11. Krishnamurphy G.T., et al. Kinetics of ^{99m}Tc-Labelled Pyrophosphate

1173237

DOCUMENT SOURCE	
Lawrence Berkeley Laboratory Archives and Records Office SCIENTISTS' PAPERS	
Records Series Title	JOHN LAWRENCE
Accession No.	434-12-00161
File Code No.	10-14-6
Carton No.	②
Folder No.	
Notes	
Found By	Karen Holmer
Date	

COPY

References:

- and Polyphosphate in Man. J Nucl Med 16: 109-115, 1975.
12. Tofe, A.J. et al. Correlation of Neoplasms with Incidence and Localization of Skeletal Metastases: An Analysis of 1,355 Diphosphonate Bone Scans J Nucl Med 16: 986-989, 1975.
 13. Holder, L.E. et al. The Camut Approach to Scintigram Interpretation- Diagnostic Aid and Teaching Method. J Nucl Med 16: 1121-1124, 1975.
 14. Witherspoon, L.R. et al. Bone Scan Patterns of Patients with Diffuse Metastatic Carcinoma of the Axial Skeleton. J Nucl Med 17: 253-257, 1976.
 15. Frankel, R.J. et al. Normal Bone Radionuclide Imaging with Diffuse Skeletal Lymphoma, Radiology 111: 365-366.
 16. Rosenthal, L. and Kaye, M. Technetium-99m Pyrophosphate Kinetics and Imaging in Metabolic Bone Disease J Nucl Med 16: 33-39, 1975.
 17. Sy, W.M. et al. "Cold" Lesions on Bone Imaging. J Nucl Med 16: 1013-1016/ 1975.
 18. Goergen, T.G. et al. "Cold" Bone Lesions: A Newly Recognized Phenomenon of Bone Imaging J Nucl Med 15: 1120-1124, 1974.
 19. Vieras, F. et al. Focal Decrease Skeletal Uptake Secondary to Metastatic Disease, Radiology 118: 121-122.
 20. Park, C.H. et al. Reliability of Renal Imaging Obtained Incidentally in ^{99m}Tc-Polyphosphate Bone Scanning. J Nucl Med 14: 534, 536, 1973.
 21. Hattner, R.S. et al. Significance of Renal Asymmetry in Bone Scans: Experience in 1795 Cases. J Nucl Med 16: 161-163, 1975.
 22. Sy, W. M. Bone Scan in Primary Hyperparathyroidism. J Nucl Med 15: 1089-1091, 1974.
 23. Sy, W.R. et al. Significance of Faint or Absent Kidney Sign on Bone Scan. J Nucl Med 16: 454-456, 1975.

1173238

DOCUMENT SOURCE Lawrence Berkeley Laboratory Archives and Records Office SCIENTISTS' PAPERS	
Records Series Title	JOHAN LAWRENCE
Accession No.	432-92-0066
File Code No.	19-112-0
Carton No.	2
Folder No.	
Notes	
Found By	Karen Holmes
Date	

COPY

References:

24. Cardi J, Bonneyville B: Diagnostic Value of Hepatic Scintillography. Arch Mal Appar Dig 51: 55-82, 1962
25. Beauchamp JH, Belanger MA, Neitznman HR: Intrahepatic Focal Lesion in Acute Viral Hepatitis. J Nucl Med 15: 356-357, 1974
26. Shirazi PH, Rayudu, VS, Fordham EW: Review of Solitary F-18 Bone Scan Lesions. Radiology 112: 369-372, 1974
27. Shirazi, PH, Raydu VS, Fordham, EW: F-13 Bone Scanning: Review of Indications and Results of 1,500 scans. Radiology 112: 361-368, 1974,
28. Isitman AT, Komaki S, Holmes RA: A Benign Uptake of Tc-99m-Polyphosphate After Radical Mastectomy. Radiology 110: 159-161, 1974.
29. Thrall, JH, Chaeed N, Geslien GE, et al: Pitfalls in Tc-99m-Polyphosphate Skeletal Imaging. Am J Roentgenol, Rad Therapy and Nuclear Med 121: 739-747, 1974.
30. Nelson L, Watson RC, Benua RS, et al: F-18 Radioisotope Scanning of Metastatic Bone Lesions in Children with Neuroblastoma. Am J Roentgenol, Rad Therapy and Nuclear Med 115: 191-199, 1972.
31. Wanken JF, Eyring EJ, Samuels LD: Diagnosis of Pediatric Bone Lesions: Correlation of Clinical, Roentgenographic, Sr-87m Scan, and Pathologic Diagnosis. J Nucl Med 14: 803-806, 1973.
32. Blau M, Ganatra R, Bender MA : F-18 Fluoride for Bone Imaging. Semin Nucl Med 2: 31-37, 1972.
33. Moon NF, Dworkin III, LaFluer PD: The Clinical Use of Sodium Fluoride F-18 in Bone Photoscanning. JAMA 204: 116-122, 1968
34. Papavasiliou C, Kostamis P, Angelakis P, et al: Localization of Sr-87m in Extrasosseous Tumor. J Nucl Med 12: 265-268, 1971
35. Samuels LD: Skeletal Scintigraphy in Children. Semin Nucl Med 2: 89-107, 1972

1173239

DOCUMENT SOURCE Lawrence Berkeley Laboratory Archives and Records Office SCIENTISTS' PAPERS	
Records Series Title	JOHN LAWRENCE
Accession No.	43a-92-016a
File Code No.	18-14-6
Carton No.	2
Folder No.	
Notes	
Found By	Karen Holmes
Date:	

COPY

References

36. Samuels LD: .Diagnosis of Malignant Bone Disease with Strontium. Can Med Assoc J 104: 411-413, 1971
37. Rosenfield N, Treves S: Osseous and Extrasosseous Uptake of Fluorine-18 and Technetium-99m-Polyphosphate in Children with Neuroblastoma. Radiology 111: 127-133, 1974
38. Fitzner PM: Tc-99m-Polyphosphate Concentration in a Neuroblastoma. J Nuc Med 15: 904-906, 1974
39. Nolan NG: Intense Uptake of Tc-99m-Diphosphonate by an Extrasosseous Neurofibroma. J Nuc Med 15: 1207-1208, 1974
40. Chaudhuri TK, Chaudhuri TK, Gulesserian HP, et al: Extrasosseous Non-Calcified Soft Tissue Uptake of Tc-99m-Polyphosphate. J Nuc Med 15: 1054-1056, 1974
41. Wenzel WW, Heast RG: Uptake of Tc-99m-Stannous Polyphosphate in an Area of Cerebral Infarction. J Nuc Med 15: 207-209, 1974
42. Chaudhuri TK, Chaudhuri TK, Suzuki Y: Splenic Accumulation of Sr-87-m in a Patient with Hodgkin's Disease. Radiology 105: 617-618, 1972
43. Suzuki Y, Hisada K, Takida M: Demonstration of Myositis Ossificans by Tc-99m-Pyrophosphate Bone Scanning. Radiology 111: 663-664, 1974
44. McQuade S, Higgins HP: Tc-99m-Polyphosphate in Diagnosing Meningioma of the Sphenoid Wing. J Nuc Med 15: 1205-1206, 1974
45. Heiser WJ, Quinn JL, Mollihan WV: The Crescent Pattern of Increased Radioactivity in Brain Scanning. Radiology 87: 483-488, 1966
46. Forster DMC, Bethell AN: The Diagnostic Value of Scintillation Brain Scanning. Clin Radiol 20: 257-268, 1969
47. Subramanian G, McAfee JG, Blair RT, Kalfelz FA, Thomas FD: Technetium 99m Methylene Diphosphonate - A Superior Agent for Skeletal Imaging: Comparison with other Technetium Compounds. J Nuc Med 16: 744-755, 1975

1173240

DOCUMENT SOURCE	
Lawrence Berkeley Laboratory Archives and Records Office SCIENTISTS' PAPERS	
Records Series Title	JOHN LAWRENCE
Accession No.	432-92-0016
File Code No.	13-14-0
Carton No.	2
Folder No.	
Notes	
Found By	Karen Holmes
Date:	

COPY

REFERENCES

48. Lisboa R, Rennie WRJ, Daniel RK: Radionuclide Evaluation of Free Vascularized Bone Graft. American Journal of Roentgenology 134: 387-388, February 1980.
49. Valdez VA and Jacobstein JG: Decreased Bone Uptake of Technetium-99m Polyphosphate in Thalassemia Major. J Nuc Med 21: 47-49, 1980.
50. Fawcett HD, McDougall IR: Bone Scan in Extraskeletal Neuroblastoma with Hot Primary and Cold Skeletal Metastases. Clinical Nuclear Medicine 5: 49-50, February 1980.
51. Moyle JW and Spies SM: Bone Scan in a Case of Amyloidosis. Clinical Nuclear Medicine 5: 51-52, February 1980.
52. Alpert LI: Pulmonary Uptake of Gallium-67 in Wegener's Granulomatosis. Clinical Nuclear Medicine 5: 53-54, February 1980.
53. Wiener SN and Kirschenbaum D: Osteoid Osteoma Presenting as Regional Osteoporosis. Clinical Nuclear Medicine 5: 68-69, February 1980.
54. Lieberman CM and Hemingway DL: Scintigraphy of Shin Splints. Clinical Nuclear Medicine 5: 31, January 1980.
55. Glassman AB and Selby JB: Another Bone Imaging Agent False-Positive: Phimosis. Clinical Nuclear Medicine 5: 34, January 1980.
56. Lyons KP and Jensen JL: Dental Lesions Causing Abnormalities on Skeletal Scintigraphy. Clinical Nuclear Medicine 4: 509-512, December 1979.
57. Goldstein HA and Treves S: Bone Scintigraphy of Osteoid Osteoma: A Clinical Review. Clinical Nuclear Medicine 3: 359-363, September 1978.
58. Sed IS, Joo KG, Baeumler GR: Multiple Manifestations of Osteolytic Lesions on Bone Imaging Letters. Journal of Nuclear Medicine 21: 896, 1980, September.
59. Dhawan VM, Turner JW, Spencer RP: Osseous and Nonosseous "Doughnut" Sign During Bone Scanning. Clinical Nuclear Medicine 5: 423, September 1980.
60. Dessi A, Alavi A, Dalinka M et al.: Role of Bone Scintigraphy in the Evaluation and Treatment of Nonunited Fractures: Concise Communication. Journal of Nuclear Medicine 21: 931-934, October 1980.
61. Spencer RP, Datu JA: Bilateral Lower Limb Uptake of Bone Scanning Agents. Seminars in Nuclear Medicine 10: 314-316, July 1980.
62. Hughes S: Radionuclides in Orthopedic Surgery. J Bone Joint Surg Br 62-B: 141-150, May 1980.

1173241

DOCUMENT SOURCE	
Lawrence Berkeley Laboratory Archives and Records Office	
SCIENTISTS' PAPERS	
Records Series Title	JOHN LAWRENCE
Accession No.	434-92-0016
File Code No.	18-14-0
Carton No.	2
Folder No.	SARNOFF, EUGENE L.
Notes	
Found By	Karen Holmes
Date	

COPY

63. Weiss PE, Jay CM, Hoffer PB et al: ^{99m}Tc -Methylene Diphosphate Bone Imaging in the Evaluation of Total Hip Prostheses. Radiology 133: 727-729, December 1979.
64. Rosenthal L, Lisbona R, Hernandez M, et al: ^{99m}Tc -PP and ^{67}Ga Imaging Following Insertion of Orthopedic Devices, Radiology 133: 717-721, December 1979.

1173242