

DOCUMENT SOURCE	
Lawrence Berkeley Laboratory Archives and Records Office	
Biology & Medicine Division	
Records Series Title	<i>Research Medicine HEALTH EFFECTS OF MARINE FISH</i>
Accession No.	<i>72-90-0203</i>
File Code No.	<i>15-7-7</i>
Carton No.	<i>(14)</i>
Folder No.	<i>Human Use Committee</i>
Notes	
Found By	<i>Karen Holmes</i>
Date	<i>1/10-1985</i>

COPY

715381

BERKELEY: DONNER LABORATORY

19 February 1981

Refer to 81RM053

Dobie Jenkins
Committee for the Protection
of Human Subjects
A & E Building
Campus

RE: CPS #80-9-15

Dear Dobie:

Enclosed is the revised version of the referenced consent form for patients involved in both our protocol ("Brain Tumor Localization and Blood Brain Barrier Studies Using Rb-82") and Dr. Fabrikant's protocol ("Investigation of Charged-Particle Beams for Radiological Neurosurgery in Treatment of CNS Disorders"). Section 2.b. of the consent form is revised so that patients are informed that they are receiving an additional exposure of radiation.

I hope this modification is satisfactory to the Committee.

Sincerely,

Chi-Kwan Yen, M.D.
Principal Investigator

CKY:mg
Enclosure

cc: LBL Human Use Committee

3006011

DOCUMENT SOURCE	
Lawrence Berkeley Laboratory Archives and Records Office	
Records Series Title	Biology & Medicine Division
Accession No.	44-90-0265
File Code No.	15-7-7
Carton No.	(A)
Folder No.	Human Use Committee
Notes	
Found By	Karen Holmes
Date	1/16-1985

COPY

Rev. 2/81
A/V

CONSENT TO ACT AS SUBJECT

FOR

TRANSVERSE SECTION AND TOMOGRAPHIC IMAGING USING RUBIDIUM-82

The diagnostic procedure of imaging the brain in transverse section is a new method for determining the health of brain tissue. The isotope, rubidium-82, goes into brain if the brain tissue is damaged. This procedure may aid a physician in evaluating the extent and location of the blood vessel abnormalities. There is little discomfort from the injection of the isotope as the radiation dose is far less than that received from angiographic or other nuclear medicine procedures. Only a few places in the world have the instrumentation and isotopes available for this new procedure.

Subject's name _____

Date _____

1. I hereby authorize _____
(name of person(s) who will perform diagnostic procedures)
and/or such qualified assistants as may be selected by him to perform the following:

Radiographs of the brain will be made using a rubidium isotope. This isotope will be injected intravenously using saline (salt water) for the infusion. I will rest comfortably on my back and may have EKG electrodes on both arms or chest during this procedure. The information from the brain will be detected by a ring of crystal detectors. A few injections might be necessary. The entire procedure will take 1 1/2 to 2 hours.

2. I understand that the procedure described in Paragraph 1 involves the following risks and discomforts:
- a. Slight discomfort associated with placement of the indwelling intravenous needle.
 - b. The radiation received in this study is in addition to the radiation associated with therapeutic irradiation. There is some risk associated with any radiation exposure. The dosage involved in these procedures is small and is equivalent to the dose from routine nuclear medicine procedures for examination of the heart, lungs, brain or kidneys. The whole-body dose from the isotope is only 0.073 rad or equivalent to 1/2 the natural background dose for one year. The kidneys receive the maximum dose which is 4.3 rads (less than that received from a comprehensive x-ray kidney study).
3. I understand the the possible benefits of the procedure(s) and/or investigation(s) are as follows:
- a. Information will be obtained about blood vessels in the brain. This type of information cannot be obtained by other methods.

3006012

DOCUMENT SOURCE
Lawrence Berkeley Laboratory
Archives and Records Office
SCIENTISTS' PAPERS

Records Series Title JOHN LAWRENCE

Accession No. 424-92-0010
 File Code No. 14-14-6
 Carton No. 2
 Folder No. EUGENE L. SAENGER
 Notes IMPO. ON BONE SEEKING
 Found By Karen Holmes
 Date:

COPY

AGENTS & BONE IMAGING. INCLUDES INFO ON OLDER "OBSOLETE" BONE SEEKING AGENTS

L.L. Saenger, M.D.
 M. Fernandez, M.D.
 E.B. Silberstein, M.D.
 R. David, M.D.
 J. Graham, D.O.
 Radioisotope Laboratory
 University of Cincinnati College of Medicine
 1981

File under Eugene Saenger
Prophylaxis

*Need to Support 1st
 Cannot need scans
 as one needs X-ray.
 Info too incomplete
 Need for quality scans
 Probe + converging
 collimators*

BONE IMAGING

I. RADIOPHARMACEUTICALS

A. ^{99m}Tc LABELLED PHOSPHATE COMPOUNDS

These include a number of compounds but the following preparations

are most frequently used:

1. ^{99m}Tc-Ethanehydroxy Diphosphonate (^{99m}TcEHDP)
2. ^{99m}Tc-Polyphosphate (^{99m}TcPoly)
3. ^{99m}Tc-Pyrophosphate (^{99m}TcPyro)
4. ^{99m}Tc-Methylene Diphosphonate (^{99m}TcMDP)
5. ^{99m}Tc-Hydroxymethylene Diphosphonate (^{99m}TcHMDP)

Dose




FIG. 1 Chemical formulas of bone seeking phosphates and diphosphonates forming complexes of ^{99m}Tc. Although these are shown as tetrasodium salts, in solution at neutral pH they are probably disodium salts.

These five preparations have been shown to have biologic differences apparently related to their chemical structures. The mechanism of bone uptake of these compounds is not perfectly elucidated but they seem to bind strongly to the bone surface by absorption ("chemisorption"). A recent report suggests that these compounds, and especially pyrophosphate, have a greater affinity for non-osteoid organic matrix than the crystal surface.

DOCUMENT SOURCE Lawrence Berkeley Laboratory Archives and Records Office <i>SCIENTISTS' PAPERS</i>	
Records Series Title <u>JOHN LAWRENCE</u>	
Accession No. <u>434-92-0066</u>	COPY
File Code No. <u>14-14-6</u>	
Carton No. <u>2</u>	
Folder No. <u>SAENGER, EUGENE L.</u>	
Notes	
Found By <u>Karen Holmes</u>	
Date	

BONE TRACING

All five agents have a high affinity for bone but they also have biological differences which seem to be related to their differences in chemical structure (4,5).

The blood clearance of the Tc-Phosphates in humans and animals appears to be best for the MDP and HMDP, followed by EHDP, Pyrophosphate, and Polyphosphate.




FIG. 2 Blood clearance of MDP in humans compared with three other ^{99m}Tc complexes and ^{99m}Tc (corrected for physical decay), assuming blood volume was 7% of body weight. P Pi indicates pyrophosphate and Poly P denotes polyphosphate.

Subramanian et al. J Nuc Med 16: 748, 1975.

The relatively slower clearance of the ^{99m}Tc-Pyrophosphate and ^{99m}Tc-Polyphosphate is explained by the stronger affinity and binding to serum proteins.

*? Also for Van...
myocard...*

DOCUMENT SOURCE Lawrence Berkeley Laboratory Archives and Records Office SCIENTISTS' PAPERS JOHN LAWRENCE	
Records Series Title	_____
Accession No.	434-92-0016
File Code No.	19-14-16
Carton No.	②
Folder No.	_____
Notes	_____
Found By	Karen Holmes
Date:	_____

COPY

BONE IMAGING

3

Overall, Tc-diphosphonate has been favored as the best agent for clinical use followed by the Tc-pyrophosphate and the polyphosphates (10,11).

As an example, some specific data are given for the labelled diphosphonate.

B. TECHNETIUM LABELLED DIPHOSPHONATES (P-C-P Bonds [Pyrophosphonates have P-O-P bonds which lead to some minor differences]).

1. Actions

- a. Increased blood flow
- b. Chemisorption
- c. Displacement of orthophosphate

2. Localization

- a. Amount and type of calcium and phosphorus present
- b. Increased vascularity of area (i.e., epiphyses)
- c. Increased surface area
- d. Increased turnover

3. Physiology and Pharmacology

- a. Optimum scan time - 2-3 hours, depending on the pharmaceutical 50% in bone - 50% excreted. Patient should void at 1 hour and at time of imaging.
- b. Diphosphonate - chelate excreted via GFR (bladder is critical organ)
- c. High ratio bone/soft tissue
- d. High ratio lesion to normal bone
- e. Low radiation dose

3006015

DOCUMENT SOURCE Lawrence Berkeley Laboratory Archives and Records Office <i>SCIENTISTS' PAPERS</i>	
Records Series Title	JOHN LAWRENCE
Accession No.	92-92-0066
File Code No.	14-14-6
Carton No.	(2)
Folder No.	
Notes	
Found By	Karen Holmes
Date:	

COPY

TABLE I

DOSIMETRY OF BONE IMAGING AGENTS

(in 70 kg Adult - Rads/mCi Administered)

(Data from package insert)

ORGAN	Radiopharmaceutical				
	^{99m} TcMDP	^{99m} Tc Pyrophosphate	^{99m} Tc EHDP	¹⁸ F	⁶⁷ Ga Citrate
Skeleton	.035	.034	.039	.15-.29	.44
Bone Marrow	.028	.028-.036	.028	.04	.58
Kidneys	.031	.028-.14	.14		.41
Liver	.008	.001	----		.46
Total Body	.0065	.009	.009	.05	.26
Bladder				2.0	
2-3 h voiding	.13	.04-.1	.1		----
4.8-6 h voiding	.31	.07-.23	.23		----
Testis					.24
2-3 h voiding	.008	.004-.01			
4.8-6 h voiding	.01	.007-.015			
Ovaries					.28
2-3 h voiding	.012	.004-.009			
4.8-6 h voiding	.017	.007-.015			
Heart					----
Normal		.007	.007		
Impaired		.014	.02-.03		
Gastroenteric Tract					
Stomach					.22
Small bowel					.36
Upper large bowel					.56
Lower large bowel					.9

Doses 10-20 mCi
Blood pool pap infection
average 3-5 hrs, 24 hrs.
35 mRads/mCi of administered activity to bone
kidneys dose 2x
marrow " 1/2
TB dose 10 mRads/mCi
Bladder wall > 10 Rads/study
hydrate + void

3006016

DOCUMENT SOURCE Lawrence Berkeley Laboratory Archives and Records Office <i>SCIENTISTS' PAPERS</i> JOHN LAWRENCE	
Records Series Title	
Accession No.	<i>92-92-0066</i>
File Code No.	<i>14-14-0</i>
Carton No.	<i>2</i>
Folder No.	
Notes	
Found By	<i>Karen Holmes</i>
Date	

COPY

BONE IMAGING

C. OLDER BONE SEEKING AGENTS

1. ^{18}F - short T_{1/2} (1.8 hr). High energy (511 KeV) annihilation photons are not ideal for gamma camera.

^{18}F has very favorable biologic characteristics. It is absorbed by the GI tract and can be given orally. The mechanism of uptake appears to be that of ion exchange at the level of the hydroxyapatite crystals. Extraction efficiency is excellent and approaches 100% depending on bone flow.

^{18}F is rapidly secreted by kidney (20-60% at 2 hours) a factor to keep in mind since overlapping of the kidneys and bladder with structures may occur.

2. ^{85}Sr - long T_{1/2} (64 days) requires low administered dose for low patient doses. This leads to low photon flux, long scanning time, and low resolution. Obsolete.

3. ^{87}Sr - short T_{1/2} (2.8 hours) does not allow for adequate soft tissue clearance. Higher photon energy (388 KeV) not ideal for gamma cameras. Obsolete.

DOCUMENT SOURCE Lawrence Berkeley Laboratory Archives and Records Office SCIENTISTS' PAPERS	
Records Series Title	<u>JOHN LAWRENCE</u>
Accession No.	<u>434-92-0216a</u>
File Code No.	<u>14-14-0</u>
Carton No.	<u>2</u>
Folder No.	
Notes	
Found By	<u>Karen Holmes</u>
Date	

COPY

BONE IMAGING

SS
SD Scan
Hodgkin

II. PATHOPHYSIOLOGY

It appears that several factors are implicated and determine the degree of concentration of bone seeking agents in the osseous tissue.

These factors are:

1. Vascularity and blood flow
2. Deposition at the level of the mineral phase (hydroxyapatite)
3. Deposition at the organic phase (osteoid matrix)
4. Degree and level between osteoblastic and osteolytic activities
5. Finally the mechanism of deposition in some non-osteoid forming tissues like tumors and ischemic and necrotic tissues (e.g. myocardial infarction), is not fully understood but intracellular metabolic changes appear to be involved.

The relationship between osteolytic and osteoblastic activities is a very important factor in determining deposition of radionuclides in bone. Radionuclide uptake is associated with osteoblastic activity. Since most of the pathologic processes are associated with osteoblastic activity (tumor, infection, trauma, etc.) bone scanning is a non-specific procedure. Final diagnosis should be attempted on other clinical grounds. One expects to find positive bone scans when osteoblastic activity is the prominent event and conversely negative studies if absent. This finding is the common one, but there are many exceptions to this observation.

III. INTERPRETATION - NORMAL

A. The Spine

Some difference due to tomographic effect of focused collimators used with rectilinear scanners versus gamma cameras must be considered.

1. Vertebral Bodies

- a. Best on posterior position where lumbar region is seen as being of uniform density. Hip and femoral heads are well seen especially with large patients.

B. The Skull

1. Cranial vault and facial structures are well seen (sutures seen in children).
2. Base of skull
3. External occipital protuberance
4. If positive image is seen, obtain vertex view to exclude intracranial disease.

3 slides
1 front
2 views
2 anterior
1 superior
1 lateral
low photon efficiency X-rays

DOCUMENT SOURCE Lawrence Berkeley Laboratory Archives and Records Office SCIENTISTS' PAPERS.	
Records Series Title	JOHN LAWRENCE
Accession No.	434-72-010a
File Code No.	18-14-6
Carton No.	(2)
Folder No.	
Notes	
Found By	Karen Holmes
Date	

COPY

BONE IMAGING

C. The Thoracic Region

1. Inferior angle and spine of scapula are well seen. Sternum and ossification centers and sternoclavicular joints (watch for assymetry. Occasionally, costo chondral junctions are well seen.

D. The Long Bones

1. Small bones in wrists, hands, and feet are frequently easily visible due to large proportion of cancellous to cortical bones. Good detail achieved with pinhole collimator.

E. Other Normal Sites of Increased Uptake

1. Epiphyses and apophyses in juveniles.
2. Joints - variable and sometimes assymetric.
 - a. Sacro-iliac joints usually show increased uptake. Sacral tubercle normally prominent (~4%). [Blei L., et al., The Sacral Tubercle - A Cause for Hot Spots on Bone Scan. Clin Nuc Med 3: 351-354, 1978].
 - b. Skull - petrous pyramids and facial bones
 - c. Cartilage - watch for thyroid cartilage
 - d. Shoulder uptake corresponding to handedness and/or position.
3. Organs
 - a. Kidney and bladder
 - b. Breast
 - c. Myocardium
4. Free ^{99m}Tc O₄
 - a. Thyroid
 - b. Salivary Gland, Sinuses
 - c. Stomach
 - d. Bowel
 - e. Liver (rare)
5. Technical - change in distance of spine from collimator

* IV. INTERPRETATION - ABNORMAL

- * A. Non-specificity of Bone Scans - cannot distinguish benign from malignant disease.

B. Soft Tissue Injury

Post-operative sites, myocardial infarction, CVA.

C. Vertebral Bodies

Most common sites for bony metastasis.

D. Delayed Scanning

- * 1. In patients who cannot empty bladder; to increase target/non-target ratio.

DOCUMENT SOURCE	
Lawrence Berkeley Laboratory Archives and Records Office	
Records Series Title <u>SCIENTISTS' PAPERS</u> <u>JOHN LAWRENCE</u>	
Accession No.	<u>434-92-0016a</u>
File Code No.	<u>14-14-6</u>
Carton No.	<u>2</u>
Folder No.	
Notes	
Found By	<u>Karen Holmes</u>
Date	

COPY

BONE IMAGING

8
Vording

D. Delayed Scanning (continued)

2. In patients with questionable densities at 2-3 hours. At 12-24 hours occasionally an abnormality will appear better visualized than at the earlier time.

E. Spot Images using pinhole or other collimator (converging or diverging) as indicated is often useful. Oblique and lateral views are often helpful. The opposite side should be imaged for comparison. *

F. Metastases

Metastases to the skeleton may show as single or multiple areas of increased uptake. Practically any bone of the skeleton may be involved, but spine, ribs, and pelvis are the most frequently observed.

The most frequent tumors with bone metastases found in clinical practice are cancer of the breast, lung and prostate gland (12).

A clear decision for preoperative bone scans in patients with operable Breast Cancer is not entirely resolved. Patients with lesions < 2 cm in diameter tend not to have positive scans. With increase in lesion size and the presence of positive axillary nodes rates of true positive scans range from 6% to about 25%. The variation is due to several factors: size of tumor, differences in hospital populations, differences in technique, problems of FP and FN interpretation. FN include early very small metastases, lack of an osteoblastic response, lesions of pubis and ischium. FP include the many other causes of positive bone scans. CT, observation and biopsy may be needed to resolve this problem. The subsequent development of metastases is related to the size and extent of the original tumor.

References: Bone scanning in breast cancer, Brit. M.J. 2: 180-181, 15 July 1978.

Galasko CSB. Problems associated with the detection of skeletal metastases. J. Roy. Soc. Med. 71: 38-41, 1978.

McNeil BJ et al. Preoperative and follow up of bone scans in patients with primary carcinoma of the breast. Surg, Gynec and Obst 147: 745-748, 1978.

Clark DG et al. Indications for bone scans in preoperative evaluation of breast cancer. Am J Surg 135: 667-670, 1978.

G. The Patient With Single Bone Lesions

Single bone lesions in bone scanning are frequently found (13). The most common causes are:

1. Metastatic tumor (lung, breast, prostate)
2. Post thoracotomy (surgery scar)
3. Fracture of vertebral body
4. Normal variants (shoulder, sternum)

3006020

DOCUMENT SOURCE Lawrence Berkeley Laboratory Archives and Records Office <i>SCIENTISTS' PAPERS</i>	
Records Series Title	<u>JOHN LAWRENCE</u>
Accession No.	<u>72-92-0016</u>
File Code No.	<u>14-14-6</u>
Carton No.	<u>2</u>
Folder No.	
Notes	
Found By	<u>Karen Holmes</u>
Date	

COPY

BONE IMAGING

H. The Patient with Diffuse Increase Uptake in Skeleton (14,15,16)

Diffuse or symmetric increase uptake is not a rare situation. The importance of this possibility stems from the fact that a symmetric diffuse uptake may be easily overlooked.

This finding has been noted in metastatic cancer of the prostate and diffuse lymphoma and cancer of the renal pelvis. It is also expected in metabolic bone disease due to hyperparathyroidism and osteomalacia. In our lab we have found Paget's of the skull and hypertrophic pulmonary osteoarthropathy present with symmetric bone involvement. It can occur in hypervitaminosis D and also in renal osteodystrophy where it may represent secondary hyperparathyroidism. [Fogelman et al. Bone scan findings in hypervitaminosis D: Case report, J Nuc Med 18: 1205-1207, 1977].

I. Presence of "Cold" Lesions in Bone Scanning (17,18,19)

A cold lesion may occur as the sole abnormality in a bone scan and it has been recognized most frequently in the vertebra and long bones. It has been described in the following conditions:

1. Metastatic disease of different sources
2. Post-traumatic aseptic necrosis
3. Sickle cell anemia with bone infarction
4. Chronic renal failure

Pathogenesis of this type of condition has not been elucidated but interruption of blood supply appears to be a prominent factor. Conceivably a poor osteoblastic reaction surrounding an area of necrosis or tumor would result in a "cold" area.

See Table III.

J. The False Negative and False Positive Bone Scan

False negative bone scans may result when radioactivity in the lesion is not intense enough, when the lesion is mostly osteolytic (e.g., 50% of multiple myeloma) or when the disease process involves the skeleton in a diffuse or symmetric fashion.

False positive scans are seen mainly as a consequence of contamination with urine and at the site of injection. If the technetium label is poor, uptake by the thyroid gland and the stomach may be a source of confusion. Increased uptake in the area of radical mastectomy presumably due to decreased attenuation of γ rays by removed soft tissue has been observed. Increased uptake in areas of post-surgery scars, biopsy sites, normal or pathologic breast tissue, soft tissue, inflammation and myocardial and cerebral infarcts occurs.

K. Increased Concentration in Extraskeletal Tissues

This condition has been found in many circumstances. The causes and conditions can be listed as follows:

1. Uptake in Osteoid Forming Tissues

3006021

DOCUMENT SOURCE	
Lawrence Berkeley Laboratory Archives and Records Office	
SCIENTISTS' PAPERS	
Records Series Title	JOHN LAWRENCE
Accession No.	424-92-0066
File Code No.	18-14-6
Carton No.	2
Folder No.	
Notes	
Found By	Karen Holmer
Date	

COPY

BONE IMAGING

10

K. Increased Concentration in Extraskelatal Tissues (con't.)

- a. Myositis ossificans
- b. Diffuse calcinosis
- c. Metastatic osteosarcoma
- d. Osseous metaplasia in soft tissues or lymph nodes
- e. Ectopic calcification in paraplegic patients
- f. Metastases of bladder (with osseous mataplasia)
- g. Calcific tendonitis

In the above conditions concentration of bone seeking agents is expected and probably related to the presence of hydroxyapatite as well as immature collagen matrix.

2. "Non-osteoid" forming tissues

- a. Thoracotomy and post-surgery scars
- b. Soft tissue inflammation
- c. Brain infarction
- d. Brain astrocytoma
- e. Brain metastases from lymphoma
- f. Myocardial infarction

The mechanism of uptake is not known but increased blood flow and concentration of the isotope by the collagen and intracellular structure could be involved.

L. The Significance of Kidney Abnormalities Incidentally Found During Bone Imaging (20-23)

This point has been recently brought to attention since many of the bone seeking radioisotopes, specifically the Tc-Phosphate compounds, are avidly concentrated by the kidneys enabling us to observe some anatomic detail of the kidneys and excretory system. The following points should be emphasized:

1. There is no good correlation between kidney concentration of ^{18}F -Fluoride and renal function.
2. Hence, kidney asymmetry during ^{18}F -Fluoride bone scanning is unreliable to predict renal disease.
3. The overall accuracy and the level of confidence when kidneys are found morphologically abnormal are high.
4. In general, information regarding kidney morphology and function appears to be similar with all four currently used Tc-Phosphate compounds.
5. The most common conditions that can be detected during bone imaging in general clinical practice are -
 - a. Bilateral decreased visualization due to:
 1. bilateral kidney disease
 2. increased bone uptake ("steal phenomena")

3006022

DOCUMENT SOURCE Lawrence Berkeley Laboratory Archives and Records Office <i>SCIENTISTS' PAPERS</i>	
Records Series Title	<u>JOHN LAWRENCE</u>
Accession No.	<u>434-92-0216a</u>
File Code No.	<u>14-14-0</u>
Carton No.	<u>2</u>
Folder No.	
Notes	
Found By	<u>Karen Holmes</u>
Date	

COPY

BONE IMAGING

11

L. The Significance of Kidney Abnormalities... (con't)

- b. Small kidney due to unilateral disease
- c. Space occupying lesions: tumors, cysts, dilated calyces
- d. Absent kidney: congenitally, surgically, destruction by encroaching process
- e. Obstructive uropathy with dilated excretory system
- f. Displacement of bladder by pelvic tumor
- g. Ectopic location

M. Bone Scanning in Pediatrics

Some pertinent points are to be considered when dealing with the pediatric age group.

1. The presence of normal variants inherent to this group such as the increased uptake in juxta-articular areas.
2. Positive bone scans due to metastatic disease of pediatric neoplasms, i.e. Wilms Tumor, neuroblastoma, leukemia, rhabdomyosarcoma. Valuable in following course of disease and evaluating treatments. In neuroblastoma bone imaging may miss the lesion although x-ray may be positive. [Reference - Kaufman RA et al. False negative bone scans in neuroblastoma metastatic to ends of long bones. Am J. Roentgenol. 130: 131-135, 1978].
3. Frequency of solitary lesions are more likely to be associated with primary bone tumors. Benign tumors take up less activity than do malignant ones. Imaging is useful in determining extent of osteoblastic tumors.
4. Differential diagnoses between osteomyelitis and cellulitis. This is a frequent situation which poses a difficult problem since clinical and radiologic findings may not be conclusive. In this regard, comparing early "blood pool" pictures with the conventional delayed scan is often helpful. In soft tissue inflammation or infection, the early blood pool scans show a greater concentration of the radioisotope when compared with the delayed scan. The reverse is usually found in cases of osteomyelitis. High sensitivity and pinhole collimators are often helpful when dealing with small children. Septic arthritis, discitis; (spondylarthritis), bone infarcts in sickle cell disease require consideration in the differential diagnosis.
5. Aseptic necrosis. If imaged early, may show decreased uptake. Later uptake will be increased. In Perthe's disease osteotomy may decrease uptake. Attention to technique important for bilateral disease and to follow course. Quantitative technique may be valuable.

[References: Norley TR et al. Femoral head activity in

3006023

DOCUMENT SOURCE Lawrence Berkeley Laboratory Archives and Records Office <i>SCIENTISTS' PAPERS</i>	
Records Series Title	<u>JOHN LAWRENCE</u>
Accession No.	<u>93A-92-0066a</u>
File Code No.	<u>19-14-6</u>
Carton No.	<u>(2)</u>
Folder No.	_____
Notes	_____
Found By	<u>Karen Holmes</u>
Date	_____

COPY

BONE IMAGING

12

M. Bone Scanning in Pediatrics (con't)

Perthe's disease: Clinical evaluation of a quantitative technique for estimating tracer uptake, J Nucl Med 19: 884-890, 1978.

Sty JR. Panner's disease (Osteonecrosis of the capitellum). Clin Nuc Med 3: 117, 1978].

6. Diaphyseal Spondylarthritis. Originally described by

DOCUMENT SOURCE Lawrence Berkeley Laboratory Archive and Records Office <i>SCIENTISTS' PAPERS</i>	
Records Series Title	<i>JOHN LAWRENCE</i>
Accession No.	<i>434-92-001616</i>
File Code No.	<i>19-14-0</i>
Carton No.	<i>(2)</i>
Folder No.	
Notes	
Found By	<i>Karen Holmes</i>
Date:	

COPY

BONE IMAGING

13

VI. GALLIUM 67 FOR BONE IMAGING

⁶⁷Ga may be useful in circumstances where ^{99m}Tc phosphorus compounds appear to give normal images as occasionally in osteomyelitis, septic arthritis. In primary bone tumors it correlates well with ^{99m}Tc compounds. It may be more useful in showing the extent of soft tissue extension. Its chief drawbacks are the high energy of its photons resulting in poor resolution. It is superior to radiography but generally not as reliable as ^{99m}Tc compounds. In selected cases where ^{99m}Tc appears equivocal or at variance with expectation it should be utilized. (Reference - Handmaker H et al. Gallium imaging in pediatrics. J Nuc Med 18: 1057-1063, 1977).

VII. DISEASES OF JOINTS

Joint abnormalities frequently show increased activity with ^{99m}Tc phosphorus compounds. Osteoarthritis joints are frequently visualized in the course of bone surveys. It is prudent to examine these joints clinically or by x-ray to exclude other causes for visualization.

In rheumatoid arthritis bone scanning agents usually show more joints to be involved than are found clinically and other forms of arthritis show similar findings. Septic arthritis is readily visualized. The peri-articular activity is usually increased due to increased blood flow and local reaction. Symmetrical abnormal increased in activity of involved joints is sometimes difficult to resolve especially with involvement of sacro-iliac joints. Although some investigators find obtaining joint to sacrum count ratios to be valuable in finding and staging sacro-iliac disease (Dequeker et al. Scintigraphic investigation of sacro-iliac disease. J Nuc Med 19: 119-120, 1978), and others (Lentle BC et al. J Nuc Med 19: 120, 1978) disagree in its usefulness for the diagnosis of ankylosing spondylitis.

A painful total hip or knee prosthesis suggests possibility of loosening of one or more components. Imagine with a Tc phosphate complex is an effective screening procedure to determine loosening or other complications. The bone scan alone lacks specificity but approaches 100% sensitivity for loosening and or infection. If the bone scan is negative, no further work-up is necessary at that time. Recent interest has been directed to the combined use of Tc-MDP and gallium in the evaluations of a painful total hip prosthesis. Both studies are needed as the interpretation depends on the presence of abnormal uptake plus the pattern of abnormal uptake, that is to assess the congruity or incongruity of the patterns of abnormal uptake.

EVALUATION OF TOTAL JOINT REPLACEMENT WITH TC-PHOSPHATES AND GALLIUM

	BONE SCAN	GALLIUM SCAN	CONGRUENT	INCONGRUENT
LOOSENING	+	-	-	-
INFECTION	+	+	-	+
OR	+	(INTENSE)	+	-
TRAUMA	+	+	+	-

3006025

VIII. QUANTITATIVE TECHNIQUES

Among these methods currently utilized is that using probes (Park et al. A quantitative evaluation of rheumatoid arthritis activity with Tc 99m HEDP, J Nuc Med 18: 973-976, 1977) with counts over affected and normal areas at specific times after:

DOCUMENT SOURCE Lawrence Berkeley Laboratory Archives and Records Office <i>SCIENTISTS' PAPERS</i> JOHN LAWRENCE	
Records Series Title	
Accession No.	93A-92-00166
File Code No.	19-14-10
Carton No.	2
Folder No.	SABINABR, EUGENE L.
Notes	
Found By	Karen Holmes
Date	

COPY

BONE IMAGING

review of earlier images to evaluate treatment in rheumatoid arthritis. Another method uses regions of interest and external standards to estimate the amounts of activity taken up in various parts of the skeleton in Paget's disease [Lurye DR et al. An improved method for quantitative bone scanning, J Nuc Med 18: 1069-1073, 1977]. A third method uses serial measurements at frequent intervals up to 24 hours with a shadow shield whole body monitor [Fogelman et al. The use of whole body retention of Tc 99m diphosphonate in the diagnosis of metabolic bone disease J Nuc Med 19: 270-275, 1978; see also Holmes RA, editorial, Quantification of skeletal Tc-99m labeled phosphates to detect metabolic bone disease, J Nuc Med 19: 330-331, 1978]. This technique differentiated patients with renal osteodystrophy, Paget's disease, osteomalacia and primary hyperparathyroidism from normal subjects. Patients with osteoporosis did not differ from normals.

IX. MISCELLANEOUS

Careful evaluation of jaws and related structures can reveal positive images associated with extraction sites, pulp and periodontal infections and local irritation [Tow DE et al. Bone scan in dental disease, J Nuc Med 19: 845-847, 1978].

The use of single photon emission tomography of facial bones using a special collimator and Tc Pyrophosphate compounds gives improved resolution of facial bones and improves the study of tumors, infections, bone grafts and post irradiation osteonecrosis [Brown ML et al. Facial bone scanning by emission tomography, J Nuc Med 18: 1184-1188, 1977].

X. GAMUT APPROACH AND REFERENCES

The gamut approach to roentgenology has been applied to nuclear medicine. Though scintiscan abnormalities are less specific than roentgen lesions, the approach has been useful.

A recent gamut prepared by one of our residents is included. It contains a comprehensive list of references.

TABLE II

SINGLE LOCALIZED AREA OF INCREASED RADIONUCLIDE UPTAKE ON A BONE SCAN

COMMON

1. Metastatic tumor (lung, breast, prostate) (26,67)
2. Post thoracotomy/surgery scar (26,28)
3. Vertebral body compression fracture (26)
4. Normal variants (shoulder, sternum) (29)

DOCUMENT SOURCE Lawrence Berkeley Laboratory Archives and Records Office SCIENTISTS' PAPERS	
Records Series Title	JOHN LAWRENCE
Accession No.	424-92-00161a
File Code No.	19-14-6
Carton No.	(2)
Folder No.	
Notes	
Found By	Karen Holmes
Date:	

COPY

BONE IMAGING

15

TABLE II (con't)

LESS COMMON

1. Metastatic tumor (cervix, neuroblastoma in children) (27,30)
2. Primary bone tumor (Ewing's sarcoma, osteosarcoma, osteochondroma) (26,31)
3. Lymphoma (26,27)
4. Monarticular degenerative disease (26)
5. Trauma - fracture other than vertebral body, biopsy site, prosthesis site) (26)
6. Osteomyelitis, pyogenic (26)
7. Peridontal disease (33) and post tooth extraction
8. Paget's disease (26,27)

RARE

1. Metastatic tumor (thyroid, renal, melanoma, pancreas, other gastrointestinal) (27)
2. Primary bone tumor (fibrosarcoma, chondrosarcoma, giant cell tumor, fibrous dysplasia, enchondroma, osteoid osteoma, bone cyst, hemangioma) (26, 32-35)
3. Multiple myeloma (27)
4. Aseptic necrosis (32)
5. Osteitis pubis (32)
6. Osteomyelitis, TB (26,36) coccidiomycosis (27)
7. Extra skeletal calcified and non-calcified tissue uptake [breast carcinoma (29,34), neuroblastoma (37,38), neurofibroma (39), brain metastasis from lung carcinoma (34), cecal/rectal carcinoma (34), nasopharyngeal carcinoma (34), fibrosarcoma (34), soft tissue abscess (40), brain infarction (41), myocardial infarction and other areas of tissue necrosis, Hodgkin's involving spleen (42), leukemic infiltrates (35), myositis ossificans (43), calcific tendonitis (32), thrombophlebitis (29)]. Also in lung, stomach, liver, kidneys in metastatic melanoma and chronic renal disease [Veukatesh et al. Metastatic calcification: the role of bone scanning, Radiology 129: 755-758, 1978; Rosenthal DI, Uptake of bone imaging agents by diffuse pulmonary metastatic calcification, Am J Roentgenol 129: 871-874, 1977; Oren VO et al., Liver metastases of oat cell carcinoma of lung detected on ^{99m}Tc diphosphonate bone scan, Clin Nuc Med 3: 355-357, 1978 - has a good table of soft tissue uptake with ^{99m}Tc bone imaging agents].

3006027

DOCUMENT SOURCE Lawrence Berkeley Laboratory Archives and Records Office <i>SCIENTISTS' PAPERS</i>	
Records Series Title	<i>JOHN LAWRENCE</i>
Accession No.	<i>424-92-0016</i>
File Code No.	<i>19-14-0</i>
Carton No.	<i>2</i>
Folder No.	
Notes	
Found By	<i>Karen Holmes</i>
Date	

COPY

BONE IMAGING

16

TABLE II (con't)

8. Gout (26)
9. Periostitis (31)
10. Meningiomas (44)
11. Scurvy - [Front D et al., Bone scintigraphy in scurvy, J Nuc Med 19: 916-917, 1978].

TABLE III

DECREASED RADIONUCLIDE UPTAKE ON A BONE SCAN

AVASCULAR AREAS

Aseptic necrosis, bone infarcts
 Radiation therapy (late effect)
 Lack of weight bearing stress
 Metastasis
 Sickle Cell C crisis
 Chronic renal failure
 Acute osteomyelitis
 References: Georgen et al., J Nuc Med 15: 1120-1124, 1974
 Fordham et al., Sem Nuc Med 4: 411-429, 1974
 Quint PA, Radiol 130: 751-752, 1979, #58.

TABLE IV

True Abnormal, False Abnormal and False Normal Abnormalities in Bone Imaging
 Silberstein, EB, J Nuc Med 17: 229-232, 1976

Abnormalities generally show increased uptake of radiopharmaceuticals

- A. True Abnormal
 1. Tumor, primary or secondary
 2. Fractures and surgical osteotomy
 3. "Metabolic"
 - a. hyperparathyroidism
 - b. Paget's disease
 - c. osteoporosis
 - d. osteomalacia and occasional pseudofractures of ribs [Fogelman et al., The role of bone scanning in osteomalacia, J Nuc Med 19: 245-248, 1978; Fogelman et al., Pseudofracture of the ribs detected by bone scanning, J Nuc Med 18: 1236-1237, 1977].
 4. Inflammation of bone
 - a. osteomyelitis
 - b. abscess
 - c. sterile osteitis (e.g. osteitis pubis)
 - d. granuloma including sarcoid, eosinophilic granuloma
 - e. hyperostosis frontalis interna
 - f. fibrous dysplasia
 - g. hypertrophic pulmonary osteoarthropathy

3006028

DOCUMENT SOURCE	
Lawrence Berkeley Laboratory Archives and Records Office SCIENTISTS' PAPERS	
Records Series Title	JOHN LAWRENCE
Accession No.	424-92-0066
File Code No.	19-14-6
Carton No.	(2)
Folder No.	
Notes	
Found By	Karen Holmes
Date:	

COPY

BONE IMAGING

17

TABLE IV (con't)

5. Arthritis
 - a. osteoarthritis
 - b. rheumatoid arthritis
 - c. gouty arthritis
 6. Soft-tissue calcifications
 - a. myositis ossificans
 - b. soft-tissue osseous metaplasia
 - c. soft-tissue tumors with calcification or ectopic bone formation
 - d. vascular calcification, especially femoral artery
 - e. calcific tendonitis
 - f. abscess
 - g. infarct, cerebral or myocardial
 - h. thrombophlebitis
 7. Vascular
 - a. surrounding the bone infarct
 8. Decreased uptake
 - a. tumor
 - b. disuse of limb (may also be increased with osteoporosis)
 - c. vascular obstruction (e.g., sickle cell disease, aseptic necrosis)
- B. Falsely Abnormal
1. Renal artifacts or disease
 - a. hydronephrosis with ^{99m}Tc -diphosphonate
 - b. contamination of clothing or skin with urine
 2. Recent surgical procedures on bone or soft tissue
 3. Biopsy site
 4. Colloid formation with liver-spleen uptake
- C. Falsely Normal
1. Lesions of smaller size than the resolving power of the system
 2. Purely lytic lesions (e.g., some myelomas)
 3. Jewelry, prostheses, pacemaker overlying a lesion

TABLE V

Bilateral Lower Limb Uptake of Bone Scanning Agents, in
R.P. Spencer and J.A. Datu, Sem Nuc Med 10, (#3): 314-316, 1980

Some causes of bilateral lower limb uptake of bone imaging agents are:

COMMON

1. "Calcified" femoral vessels
2. Trauma (and battered child)
3. Stress fractures
4. Aseptic necrosis,* osteonecrosis

*increased uptake is likely outside of the affected region.

3006029

DOCUMENT SOURCE	
Lawrence Berkeley Laboratory Archive and Records Office SCIENTISTS' PAPERS	
Records Series Title	JOHN LAWRENCE
Accession No.	434-32-0066a
File Code No.	14-14-6
Carton No.	2
Folder No.	
Notes	
Found By	Karen Holmes
Date:	

COPY

BONE IMAGING

18

TABLE V (con't).

5. Multiple small vessel occlusion, as in SS disease*
6. Osteomyelitis (acute or chronic)
7. Hypertrophic pulmonary osteoarthropathy
8. Increased collagen turnover**
9. Paget's disease
10. Arthritides (osteoarthritis, rheumatoid)
11. Multiple metastases[†]

UNCOMMON

1. Ectopic calcifications
2. Injection sites
3. Polymyositis, myolysis, dermatomyositis
4. Soft tissue infection, infarction
5. Surgically induced (bone grafts, pinning, prostheses)
6. Septic or other emboli*
7. Bilateral sympathectomy
8. Lymphoma
9. Multiple myeloma
10. Gaucher's disease
11. Sarcoidosis^{††}
12. Scurvy
13. Myelofibrosis
14. Joint disorders

*increased uptake is likely outside of the affected region

**Includes hyperthyroidism, acromegaly, thyroid acropachy, hyperparathyroidism, renal osteodystrophy.

[†]Multiple primary bone tumors can occur but are quite rare.

^{††}Well documented radiographically, less well established by bone scans.

3006030

DOCUMENT SOURCE	
Lawrence Berkeley Laboratory Archives and Records Office	
SCIENTISTS' PAPERS JOHN LAWRENCE	
Records Series Title	
Accession No.	434-92-0010
File Code No.	10-14-0
Carton No.	2
Folder No.	
Notes	
Found By	Karen Holmes
Date	

COPY

REFERENCES

1. Blau, Monte, et al. ^{18}F -Fluoride for Bone Imaging. Sem Nuc Med Vol 2, January 1972.
2. Moon, N.M. et al. The Clinical Use of Sodium Fluoride F-18 in Bone Photoscanning. JAMA 204: 116-122.
3. O'Mara, R.E. Bone Scanning in Osseous Metastatic Disease. JAMA 229: 1915-1917.
4. Subramanian, G. et al. An Evaluation of $^{99\text{m}}\text{Tc}$ -labelled Phosphate Compounds as Bone-Imaging Agents, in Subramanian, G. et al (eds) Radiopharmaceuticals, New York, Society of Nuclear Medicine, Inc., pp 319-328, 1975.
5. Ackerhalt, R.E., et al. A Comparative Study of Three $^{99\text{m}}\text{Tc}$ -labelled Phosphorus Compounds and ^{18}F -Fluoride for Skeletal Imaging. J Nuc Med 15: 1153-1157, 1974.
6. Garcia, Daniel A., et al. Relative Accretion of $^{99\text{m}}\text{Tc}$ -Polyphosphate by Forming and Resorbing Bone Systems in Rats: Its Significance in the Pathologic Basis of Bone Scanning, J Nucl Med 17: 93-97, 1976.
7. Bowen, B.M. et al. Analysis of the Relationship between $^{99\text{m}}\text{Tc}$ -Sn-Polyphosphate and $^{99\text{m}}\text{Tc}$ -Pyrophosphate. J Nucl Med 15: 652-655, 1974.
8. Rosenthal, L. and Kaye, M. $^{99\text{m}}\text{Tc}$ -Pyrophosphate Kinetics and Imaging in Metabolic Disease. J Nucl Med 16: 33, 1975.
9. Kaye, M. et al. $^{99\text{m}}\text{Tc}$ -Pyrophosphate: Studies in-vivo and in-vitro J Nucl Med 16: 40-45, 1975.
10. Krishnamurphy, G.H. et al. Clinical Comparison of the Kinetics of $^{99\text{m}}\text{Tc}$ -Labelled Polyphosphate and Diphosphonate. J Nucl Med 15: 848-855, 1974.
11. Krishnamurphy G.T., et al. Kinetics of $^{99\text{m}}\text{Tc}$ -Labelled Pyrophosphate

3006031

DOCUMENT SOURCE Lawrence Berkeley Laboratory Archives and Records Office SCIENTISTS' PAPERS JOHN LAWRENCE	
Records Series Title	
Accession No.	932-92-00166
File Code No.	14-14-0
Carton No.	2
Folder No.	
Notes	
Found By	Karen Holmes
Date:	

COPY

References:

- and Polyphosphate in Man. J Nucl Med 16: 109-115, 1975.
12. Tofe, A.J. et al. Correlation of Neoplasms with Incidence and Localization of Skeletal Metastases: An Analysis of 1,355 Diphosphonate Bone Scans J Nucl Med 16: 986-989, 1975.
 13. Holder, L.E. et al. The Gamut Approach to Scintigram Interpretation-Diagnostic Aid and Teaching Method. J Nucl Med 16: 1121-1124, 1975.
 14. Witherspoon, L.R. et al. Bone Scan Patterns of Patients with Diffuse Metastatic Carcinoma of the Axial Skeleton. J Nucl Med 17: 253-257, 1976.
 15. Frankel, R.J. et al. Normal Bone Radionuclide Imaging with Diffuse Skeletal Lymphoma, Radiology 111: 365-366.
 16. Rosenthal, L. and Kaye, M. Technetium-99m Pyrophosphate Kinetics and Imaging in Metabolic Bone Disease J Nucl Med 16: 33-39, 1975.
 17. Sy, W.M. et al. "Cold" Lesions on Bone Imaging. J Nucl Med 16: 1013-1016/ 1975.
 18. Goergen, T.G. et al. "Cold" Bone Lesions: A Newly Recognized Phenomenon of Bone Imaging J Nucl Med 15: 1120-1124, 1974.
 19. Vieras, F. et al. Focal Decrease Skeletal Uptake Secondary to Metastatic Disease, Radiology 118: 121-122.
 20. Park, C.H. et al. Reability of Renal Imaging Obtained Incidentally in ^{99m}Tc-Polyphosphate Bone Scanning. J Nucl Med 14: 534,536, 1973.
 21. Hattner, R.S. et al. Significance of Renal Asymmetry in Bone Scans: Experience in 1795 Cases. J Nucl Med 16: 161-163, 1975.
 22. Sy, W. M. Bone Scan in Primary Hyperparathyroidism. J Nucl Med 15: 1089-1091, 1974.
 23. Sy, W.R. et al. Significance of Faint or Absent Kidney Sign on Bone Scan. J Nucl Med 16: 454-456, 1975.

DOCUMENT SOURCE	
Lawrence Berkeley Laboratory Archives and Records Office SCIENTISTS' PAPERS	
Records Series Title	JOHN LAWRENCE
Accession No.	43A-92-0016a
File Code No.	19-14-12
Carton No.	(2)
Folder No.	
Notes	
Found By	Karen Holmes
Date	

COPY

References:

24. Cardi J, Bonneyville B: Diagnostic Value of Hepatic Scintillography. Arch Mal Appar Dig SI: 55-82, 1962
25. Beauchamp JH, Belanger MA, Neitzchman HR: Intrahepatic Focal Lesion in Acute Viral Hepatitis. J Nucl Med 15: 356-357, 1974
26. Shirazi PH; Rayudu, VS, Fordham EW: Review of Solitary F-18 Bone Scan Lesions. Radiology 112: 369-372, 1974
27. Shirazi, PH, Raydu VS, Fordham, EW: F-13 Bone Scanning: Review of Indications and Results of 1,500 scans. Radiology 112: 361-368, 1974.
28. Isitman AT, Komaki S, Holmes RA: A Benign Uptake of Tc-99m-Polyphosphate After Radical Mastectomy. Radiology 110: 159-161, 1974.
29. Thrall, JH, Chae N, Geslien GE, et al: Pitfalls in Tc-99m-Polyphosphate Skeletal Imaging. Am J Roentgenol, Rad Therapy and Nuclear Med 121: 739-747, 1974.
30. Nelson L, Watson RC, Benua RS, et al: F-18 Radioisotope Scanning of Metastatic Bone Lesions in Children with Neuroblastoma. Am J Roentgenol, Rad Therapy and Nuclear Med 115: 191-199, 1972.
31. Wanken JF, Eyring EJ, Samuels LD: Diagnosis of Pediatric Bone Lesions: Correlation of Clinical, Roentgenographic, Sr-87m Scan, and Pathologic Diagnosis. J Nucl Med 14: 803-806, 1973.
32. Blau M, Ganatra R, Bender MA : F-18 Fluoride for Bone Imaging. Semin Nucl Med 2: 31-37, 1972.
33. Moon NF, Dworkin HJ, LaFluer PD: The Clinical Use of Sodium Fluoride F-18 in Bone Photoscanning. JAMA 204: 116-122, 1968
34. Papavasiliou C, Kostamis P, Angelakis P, et al: Localization of Sr-87m in Extraosseous Tumor. J Nucl Med 12: 265-268, 1971
35. Samuels LD: Skeletal Scintigraphy in Children. Semin Nucl Med 2: 89-107, 1972

DOCUMENT SOURCE Lawrence Berkeley Laboratory Archives and Records Office SCIENTISTS' PAPERS JOHN LAWRENCE	
Records Series Title	
Accession No.	432-92-0066
File Code No.	19-14-6
Carton No.	(2)
Folder No.	
Notes	
Found By	Karen Holmer
Date	

COPY

References

36. Samuels LD: .Diagnosis of Malignant Bone Disease with Strontium. Can Med Assoc J 104: 411-413, 1971
37. Rosenfield N, Treves S: Osseous and Extraosseous Uptake of Fluorine-18 and Technetium-99m-Polyphosphate in Children with Neuroblastoma. Radiology 111: 127-133, 1974
38. Fitzner PM: Tc-99m-Polyphosphate Concentration in a Neuroblastoma. J Nuc Med 15: 904-906, 1974
39. Nolan NG: Intense Uptake of Tc-99m-Diphosphonate by an Extraosseous Neurofibroma. J Nuc Med 15: 1207-1208, 1974
40. Chaudhuri TK, Chaudhuri TK, Gulesserian HP, et al: Extraosseous Non-Calciified Soft Tissue Uptake of Tc-99m-Polyphosphate. J Nuc Med 15: 1054-1056, 1974
41. Wenzel WW, Heast RG: Uptake of Tc-99m-Stannous Polyphosphate in an Area of Cerebral Infarction. J Nuc Med 15: 207-209, 1974
42. Chaudhuri TK, Chaudhuri TK, Suzuki Y: Splenic Accumulation of Sr-87-m in a Patient with Hodgkin's Disease. Radiology 105: 617-618, 1972
43. Suzuki Y, Hisada K, Takida M: Demonstration of Myositis Ossificans by Tc-99m-Pyrophosphate Bone Scanning. Radiology 111: 663-664, 1974
44. McQuade S, Higgens HP: Tc-99m-Polyphosphate in Diagnosing Meningioma of the Sphenoid Wing. J Nuc Med 15: 1205-1206, 1974
45. Heiser WJ, Quinn JL, Mollihan WV: The Crescent Pattern of Increased Radioactivity in Brain Scanning. Radiology 87: 483-488, 1966
46. Forster DMC, Bethell AN: The Diagnostic Value of Scintillation Brain Scanning. Clin Radiol 20: 257-268, 1969
47. Subramanian G, McAfee JG, Blair RT, Kallfelz FA, Thomas FD: Technetium 99m Methylene Diphosphonate - A Superior Agent for Skeletal Imaging: Comparison with other Technetium Compounds. J Nuc Med 16: 744-755, 1975

4E09034

DOCUMENT SOURCE	
Lawrence Berkeley Laboratory Archives and Records Office	
SCIENTISTS' PAPERS	
Records Series Title	JOHN LAWRENCE
Accession No.	432-92-0266a
File Code No.	18-14-0
Carton No.	2
Folder No.	
Notes	
Found By	Karen Holmes
Date	

COPY

REFERENCES

48. Lisboa R, Rennie WRJ, Daniel RK: Radionuclida Evaluation of Free Vascularized Bone Graft. American Journal of Roentgenology 134: 387-388, February 1980.
49. Valdez VA and Jacobstein JG: Decreased Bone Uptake of Technetium-99m Polyphosphate in Thalassemia Major. J Nuc Med 21: 47-49, 1980.
50. Fawcett HD, McDougall IR: Bone Scan in Extraskeletal Neuroblastoma with Hot Primary and Cold Skeletal Metastases. Clinical Nuclear Medicine 5: 49-50, February 1980.
51. Moyle JW and Spies SM: Bone Scan in a Case of Amyloidosis. Clinical Nuclear Medicine 5: 51-52, February 1980.
52. Alpert LI: Pulmonary Uptake of Gallium-67 in Wegener's Granulomatosis. Clinical Nuclear Medicine 5: 53-54, February 1980.
53. Wiener SN and Kirschenbaum D: Osteoid Osteoma Presenting as Regional Osteoporosis. Clinical Nuclear Medicine 5: 68-69, February 1980.
54. Lieberman CM and Hemingway DL: Scintigraphy of Shin Splints. Clinical Nuclear Medicine 5: 31, January 1980.
55. Glassman AB and Selby JB: Another Bone Imaging Agent False-Positive: Phimosi. Clinical Nuclear Medicine 5: 34, January 1980.
56. Lyons KP and Jensen JL: Dental Lesions Causing Abnormalities on Skeletal Scintigraphy. Clinical Nuclear Medicine 4: 509-512, December 1979.
57. Goldstein HA and Treves S: Bone Scintigraphy of Osteoid Osteoma: A Clinical Review. Clinical Nuclear Medicine 3: 359-363, September 1978.
58. Sed IS, Joo KG, Baeumler GR: Multiple Manifestations of Osteolytic Lesions on Bone Imaging Letters. Journal of Nuclear Medicine 21: 896, 1980, September.
59. Dhawan VM, Turner JW, Spencer RP: Osseous and Nonosseous "Doughnut" Sign During Bone Scanning. Clinical Nuclear Medicine 5: 423, September 1980.
60. Desai A, Alavi A, Dalinka M et al.: Role of Bone Scintigraphy in the Evaluation and Treatment of Nonunited Fractures: Concise Communication. Journal of Nuclear Medicine 21: 931-934, October 1980.
61. Spencer RP, Datu JA: Bilateral Lower Limb Uptake of Bone Scanning Agents. Seminars in Nuclear Medicine 10: 314-316, July 1980.
62. Hughes S: Radionuclides in Orthopedic Surgery. J Bone Joint Surg Br 62-B: 141-150, May 1980.

3006035

DOCUMENT SOURCE	
Lawrence Berkeley Laboratory Archives and Records Office	
SCIENTISTS' PAPERS	
Records Series Title	JOHN LAWRENCE
Accession No.	434-92-0016
File Code No.	18-14-6
Carton No.	2
Folder No.	SCHNIGER, EVERETT L.
Notes	
Found By	Karen Holmes
Date	

COPY

63. Weiss PE, Jay CM, Hoffer PB et al: ^{99m}Tc -Methylene Diphosphate Bone Imaging in the Evaluation of Total Hip Prostheses, Radiology 133: 727-729, December 1979.
64. Rosenthal L, Lisbona R, Hernandez M, et al: ^{99m}Tc -PP and ^{67}Ga Imaging Following Insertion of Orthopedic Devices, Radiology 133: 717-721, December 1979.